CHLAA

Test Booklet Code

This Booklet contains 24 pages.



Do not open this Test Booklet until you are asked to do so.

Read carefully the Instructions on the Back Cover of this Test Booklet.

Important Instructions :

- 1. The Answer Sheet is inside this Test Booklet. When you are directed to open the Test Booklet, take out the Answer Sheet and fill in the particulars on **Side-1** and **Side-2** carefully with **blue/black** ball point pen only.
- 2. The test is of **3 hours** duration and this Test Booklet contains **180** questions. Each question carries **4** marks. For each correct response, the candidate will get **4** marks. For each incorrect response, **one mark** will be deducted from the total scores. The maximum marks are 720.
- 3. Use **Blue/Black Ball Point Pen only** for writing particulars on this page/marking responses.
- 4. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
- 5. On completion of the test, the candidate must hand over the Answer Sheet to the Invigilator before leaving the Room/Hall. The candidates are allowed to take away this Test Booklet with them.
- 6. The CODE for this Booklet is **HH**. Make sure that the CODE printed on **Side-2** of the Answer Sheet is the same as that on this Test Booklet. In case of discrepancy, the candidate should immediately report the matter to the Invigilator for replacement of both the Test Booklet and the Answer Sheet.
- 7. The candidates should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your Roll No. anywhere else except in the specified space in the Test Booklet/Answer Sheet.
- 8. Use of white fluid for correction is *not* permissible on the Answer Sheet.

| Name of the Candidate (in Capitals) : | |
|---|-------------------------|
| Roll Number : in figures | |
| : in words | |
| Centre of Examination (in Capitals) : | |
| Candidate's Signature : In | vigilator's Signature : |
| Facsimile signature stamp of Centre Superintendent : | |

CHLAA/HH/Page 1 Downloaded from www.cclchapter.com

- Current sensitivity of a moving coil galvanometer is 5 div/mA and its voltage sensitivity (angular deflection per unit voltage applied) is 20 div/V. The resistance of the galvanometer is
 - (1) 250 Ω
 - $(2) \quad 40 \ \Omega$
 - $(3) \quad 500 \ \Omega$
 - $(4) \quad 25 \ \Omega$
- 2. A metallic rod of mass per unit length 0.5 kg m⁻¹ is lying horizontally on a smooth inclined plane which makes an angle of 30° with the horizontal. The rod is not allowed to slide down by flowing a current through it when a magnetic field of induction 0.25 T is acting on it in the vertical direction. The current flowing in the rod to keep it stationary is
 - (1) 14·76 A
 - (2) 7.14 A
 - (3) 11·32 A
 - (4) 5·98 A
- 3. An inductor 20 mH, a capacitor 100 μ F and a resistor 50 Ω are connected in series across a source of emf, V = 10 sin 314 t. The power loss in the circuit is
 - $(1) \quad 2{\cdot}74 \ W$
 - $(2) \quad 0.79 \ W$
 - (3) 1.13 W
 - (4) 0.43 W
- A thin diamagnetic rod is placed vertically between the poles of an electromagnet. When the current in the electromagnet is switched on, then the diamagnetic rod is pushed up, out of the horizontal magnetic field. Hence the rod gains gravitational potential energy. The work required to do this comes from

 - (2) the current source
 - (3) the induced electric field due to the changing magnetic field
 - (4) the magnetic field

- The refractive index of the material of a prism is $\sqrt{2}$ and the angle of the prism is 30°. One of the two refracting surfaces of the prism is made a mirror inwards, by silver coating. A beam of monochromatic light entering the prism from the other face will retrace its path (after reflection from the silvered surface) if its angle of incidence on the prism is
- (1) 30°
- (2) 60°
- (3) zero
- (4) 45°
- An em wave is propagating in a medium with a velocity $\overrightarrow{V} = V \overrightarrow{i}$. The instantaneous oscillating electric field of this em wave is along +y axis. Then the direction of oscillating magnetic field of the em wave will be along
 - (1) -y direction
 - (2) -z direction
 - (3) -x direction
 - (4) + z direction
- 7. An object is placed at a distance of 40 cm from a concave mirror of focal length 15 cm. If the object is displaced through a distance of 20 cm towards the mirror, the displacement of the image will be
 - (1) 30 cm towards the mirror
 - (2) 30 cm away from the mirror
 - (3) 36 cm towards the mirror
 - (4) 36 cm away from the mirror
 - The magnetic potential energy stored in a certain inductor is 25 mJ, when the current in the inductor is 60 mA. This inductor is of inductance
 - $(1) \quad 1{\cdot}389 \ H$
 - (2) 0.138 H
 - (3) 13·89 H
 - (4) 138·88 H

9. An electron of mass m with an initial velocity $\overrightarrow{V} = V_0 \stackrel{\land}{i} (V_0 > 0)$ enters an electric field $\overrightarrow{E} = -E_0 \stackrel{\land}{i} (E_0 = \text{constant} > 0)$ at t = 0. If λ_0 is its de-Broglie wavelength initially, then its de-Broglie wavelength at time t is

(1)
$$\lambda_0 t$$

$$(2) \qquad \frac{\lambda_0}{\left(1+\frac{eE_0}{mV_0}t\right)}$$

$$\begin{array}{ll} (3) & \lambda_0 \\ (4) & \lambda_0 \left(1 + \frac{eE_0}{mV_0} t \right) \end{array}$$

- 10. When the light of frequency $2v_0$ (where v_0 is threshold frequency), is incident on a metal plate, the maximum velocity of electrons emitted is v_1 . When the frequency of the incident radiation is increased to $5v_0$, the maximum velocity of electrons emitted from the same plate is v_2 . The ratio of v_1 to v_2 is
 - (1) 4:1
 - (2) 1:2
 - (3) 2:1
 - (4) 1:4
- 11. For a radioactive material, half-life is 10 minutes. If initially there are 600 number of nuclei, the time taken (in minutes) for the disintegration of 450 nuclei is
 - (1) 30
 - (2) 20
 - (3) 15
 - (4) 10
- 12. The ratio of kinetic energy to the total energy of an electron in a Bohr orbit of the hydrogen atom, is
 - (1) 2:-1
 - (2) 1:1
 - (3) 1:-2
 - (3) 1:-2(4) 1:-1

Unpolarised light is incident from air on a plane surface of a material of refractive index ' μ '. At a particular angle of incidence 'i', it is found that the reflected and refracted rays are perpendicular to each other. Which of the following options is correct for this situation ?

(1)
$$i = \sin^{-1}\left(\frac{1}{\mu}\right)$$

- (2) Reflected light is polarised with its electric vector parallel to the plane of incidence
- (3) $i = \tan^{-1}\left(\frac{1}{\mu}\right)$
- (4) Reflected light is polarised with its electric vector perpendicular to the plane of incidence
- 14. In Young's double slit experiment the separation d between the slits is 2 mm, the wavelength λ of the light used is 5896 Å and distance D between the screen and slits is 100 cm. It is found that the angular width of the fringes is 0.20°. To increase the fringe angular width to 0.21° (with same λ and D) the separation between the slits needs to be changed to
 - $(1) \quad 2{\cdot}1 \text{ mm}$
 - (2) 1·8 mm
 - (3) 1·7 mm
 - (4) 1·9 mm
- **15.** An astronomical refracting telescope will have large angular magnification and high angular resolution, when it has an objective lens of
 - (1) large focal length and large diameter
 - (2) small focal length and large diameter
 - (3) small focal length and small diameter
 - (4) large focal length and small diameter


16. The power radiated by a black body is P and it radiates maximum energy at wavelength, λ_0 . If the temperature of the black body is now changed so that it radiates maximum energy at wavelength $\frac{3}{4}\lambda_0$, the power radiated by it becomes nP. The value of n is

(1)
$$\frac{256}{81}$$

(2) $\frac{3}{4}$
(3) $\frac{81}{256}$
(4) $\frac{4}{2}$

- 3
- 17. Two wires are made of the same material and have the same volume. The first wire has cross-sectional area A and the second wire has cross-sectional area 3A. If the length of the first wire is increased by Δl on applying a force F, how much force is needed to stretch the second wire by the same amount ?
 - (1) 4 F
 - (2) 9 F
 - (3) F
 - (4) 6 F
- **18.** A small sphere of radius 'r' falls from rest in a viscous liquid. As a result, heat is produced due to viscous force. The rate of production of heat when the sphere attains its terminal velocity, is proportional to
 - $(1) r^5$
 - (2) r^3
 - (3) r^4
 - (4) r^2
- 19. A sample of 0.1 g of water at 100°C and normal pressure $(1.013 \times 10^5 \text{ Nm}^{-2})$ requires 54 cal of heat energy to convert to steam at 100°C. If the volume of the steam produced is 167.1 cc, the change in internal energy of the sample, is
 - (1) 42.2 J
 - (2) 104·3 J
 - (3) 84.5 J
 - (4) 208.7 J

The kinetic energies of a planet in an elliptical orbit about the Sun, at positions A, B and C are K_A , K_B and K_C , respectively. AC is the major axis and SB is perpendicular to AC at the position of the Sun S as shown in the figure. Then




- **21.** A solid sphere is in rolling motion. In rolling motion a body possesses translational kinetic energy (K_t) as well as rotational kinetic energy (K_r) simultaneously. The ratio $K_t : (K_t + K_r)$ for the sphere is
 - (1) 10:7
 - (2) 7:10
 - (3) 2:5
 - (4) 5:7
- **22.** A solid sphere is rotating freely about its symmetry axis in free space. The radius of the sphere is increased keeping its mass same. Which of the following physical quantities would remain constant for the sphere ?
 - (1) Rotational kinetic energy
 - (2) Angular velocity
 - (3) Angular momentum
 - (4) Moment of inertia
- **23.** If the mass of the Sun were ten times smaller and the universal gravitational constant were ten times larger in magnitude, which of the following is *not* correct ?
 - (1) Time period of a simple pendulum on the Earth would decrease.
 - (2) Raindrops will fall faster.
 - (3) 'g' on the Earth will not change.
 - (4) Walking on the ground would become more difficult.

24. The moment of the force, $\overrightarrow{F} = 4\hat{i} + 5\hat{j} - 6\hat{k}$ at 28. (2, 0, -3), about the point (2, -2, -2), is given by

(1)
$$-7\hat{i} - 8\hat{j} - 4\hat{k}$$

(2) $-8\hat{i} - 4\hat{j} - 7\hat{k}$

- (3) $-7\hat{i} 4\hat{j} 8\hat{k}$
- (4) $-4\hat{i} \hat{j} 8\hat{k}$
- 25. A student measured the diameter of a small steel ball using a screw gauge of least count 0.001 cm. The main scale reading is 5 mm and zero of circular scale division coincides with 25 divisions above the reference level. If screw gauge has a zero error of 0.004 cm, the correct diameter of the ball is
 - (1) 0.053 cm
 - $(2) \quad 0{\cdot}521~\text{cm}$
 - (3) 0.529 cm
 - $(4) \quad 0{\cdot}525~cm$
- 26. A block of mass m is placed on a smooth inclined wedge ABC of inclination θ as shown in the figure. The wedge is given an acceleration 'a' towards the right. The relation between a and θ for the block to remain stationary on the wedge is



(2)
$$a = \frac{g}{cosec}$$

(3)
$$a = g \tan \theta$$

(4)
$$a = \frac{g}{\sin \theta}$$

- 27. A toy car with charge q moves on a frictionless horizontal plane surface under the influence of a uniform electric field E. Due to the force q E, its velocity increases from 0 to 6 m/s in one second duration. At that instant the direction of the field is reversed. The car continues to move for two more seconds under the influence of this field. The average velocity and the average speed of the toy car between 0 to 3 seconds are respectively
 - (1) 1 m/s, 3.5 m/s
 - (2) 2 m/s, 4 m/s
 - (3) 1.5 m/s, 3 m/s
 - (4) 1 m/s, 3 m/s

A tuning fork is used to produce resonance in a glass tube. The length of the air column in this tube can be adjusted by a variable piston. At room temperature of 27° C two successive resonances are produced at 20 cm and 73 cm of column length. If the frequency of the tuning fork is 320 Hz, the velocity of sound in air at 27° C is


- (1) 350 m/s
- (2) 330 m/s
- (3) 300 m/s
- (4) 339 m/s
- The electrostatic force between the metal plates of an isolated parallel plate capacitor C having a charge Q and area A, is
 - (1) proportional to the square root of the distance between the plates.
 - (2) independent of the distance between the plates.
 - (3) inversely proportional to the distance between the plates.
 - (4) linearly proportional to the distance between the plates.
- **30.** An electron falls from rest through a vertical distance h in a uniform and vertically upward directed electric field E. The direction of electric field is now reversed, keeping its magnitude the same. A proton is allowed to fall from rest in it through the same vertical distance h. The time of fall of the electron, in comparison to the time of fall of the proton is
 - (1) 10 times greater
 - (2) smaller
 - (3) equal
 - (4) 5 times greater
 - A pendulum is hung from the roof of a sufficiently high building and is moving freely to and fro like a simple harmonic oscillator. The acceleration of the bob of the pendulum is 20 m/s^2 at a distance of 5 m from the mean position. The time period of oscillation is
 - (1) 2 s
 - (2) $2\pi s$
 - (3) 1 s

 πs


(4)


CHLAA/HH/Page 5

32. A battery consists of a variable number 'n' of identical cells (having internal resistance 'r' each) which are connected in series. The terminals of the battery are short-circuited and the current I is measured. Which of the graphs shows the correct relationship between I and n?










- 33. A carbon resistor of $(47 \pm 4.7) \text{ k}\Omega$ is to be marked with rings of different colours for its identification. The colour code sequence will be
 - $(1) \quad Yellow-\ Green-Violet-Gold$
 - (2) Violet Yellow Orange Silver
 - (3) Green Orange Violet Gold
 - (4) Yellow Violet Orange Silver
- 34. A set of 'n' equal resistors, of value 'R' each, are connected in series to a battery of emf 'E' and internal resistance 'R'. The current drawn is I. Now, the 'n' resistors are connected in parallel to the same battery. Then the current drawn from battery becomes 10 I. The value of 'n' is
 - (1) 20
 - (2) 10
 - (3) 9
 - (4) 11

At what temperature will the rms speed of oxygen molecules become just sufficient for escaping from the Earth's atmosphere ? (Given :


Mass of oxygen molecule (m) = 2.76×10^{-26} kg Boltzmann's constant k_B = 1.38×10^{-23} J K⁻¹)

- $(1) \quad 5{\cdot}016\times 10^4 \ K$
- $(2) \quad 2{\cdot}508 \times 10^4 \; \mathrm{K}$
- $(3) \quad 1{\cdot}254\times 10^4 \; \mathrm{K}$
- (4) $8.360 \times 10^4 \text{ K}$
- **36.** The volume (V) of a monatomic gas varies with its temperature (T), as shown in the graph. The ratio of work done by the gas, to the heat absorbed by it, when it undergoes a change from state A to state B, is




- **37.** The fundamental frequency in an open organ pipe is equal to the third harmonic of a closed organ pipe. If the length of the closed organ pipe is 20 cm, the length of the open organ pipe is
 - (1) 12.5 cm
 - (2) 13.2 cm
 - (3) 16 cm
 - (4) 8 cm
- **38.** The efficiency of an ideal heat engine working between the freezing point and boiling point of water, is
 - (1) 6.25%
 - (2) 26.8%
 - (3) 12.5%
 - (4) 20%

- 39. A moving block having mass m, collides with another stationary block having mass 4m. The lighter block comes to rest after collision. When the initial velocity of the lighter block is v, then the value of coefficient of restitution (e) will be
 - (1) 0.8
 - (2) 0.5
 - (3) 0.4
 - (4) 0.25
- **40.** A body initially at rest and sliding along a frictionless track from a height h (as shown in the figure) just completes a vertical circle of diameter AB = D. The height h is equal to




- 41. Three objects, A : (a solid sphere), B : (a thin circular disk) and C : (a circular ring), each have the same mass M and radius R. They all spin with the same angular speed ω about their own symmetry axes. The amounts of work (W) required to bring them to rest, would satisfy the relation
 - (1) $W_B > W_A > W_C$
 - $(2) \quad \mathrm{W}_\mathrm{C} > \mathrm{W}_\mathrm{B} > \mathrm{W}_\mathrm{A}$
 - $(3) \quad \mathrm{W}_\mathrm{A} > \mathrm{W}_\mathrm{C} > \mathrm{W}_\mathrm{B}$
 - $(4) \quad \mathrm{W}_\mathrm{A} > \mathrm{W}_\mathrm{B} > \mathrm{W}_\mathrm{C}$
- **42.** Which one of the following statements is *incorrect* ?
 - $(1) \quad \ \ {\rm Frictional\ force\ opposes\ the\ relative\ motion.}$
 - (2) Rolling friction is smaller than sliding friction.
 - (3) Coefficient of sliding friction has dimensions of length.
 - (4) Limiting value of static friction is directly proportional to normal reaction.

In the combination of the following gates the output Y can be written in terms of inputs A and B as



- $(3) \quad \mathbf{A} + \mathbf{B}$
- (4) $\mathbf{A} \cdot \mathbf{\overline{B}} + \mathbf{\overline{A}} \cdot \mathbf{B}$
- 44. In the circuit shown in the figure, the input voltage V_i is 20 V, $V_{BE} = 0$ and $V_{CE} = 0$. The values of I_B , I_C and β are given by



- (1) $I_B = 20 \ \mu A, \ I_C = 5 \ mA, \ \beta = 250$
- (2) $I_B = 40 \ \mu A, \ I_C = 10 \ mA, \ \beta = 250$
- (3) $I_B = 40 \ \mu A, \ I_C = 5 \ mA, \ \beta = 125$
- (4) $I_B = 25 \ \mu A$, $I_C = 5 \ mA$, $\beta = 200$
- **45.** In a p-n junction diode, change in temperature due to heating
 - (1) does not affect resistance of p-n junction
 - $(2) \quad affects \ only \ reverse \ resistance$

 - (4) affects only forward resistance

Downloadest4F6FARwave.workhapter.com

| 46. | Colu | ımn Il | | | | umn I with those in <i>orrect</i> option given | 48. | Which of the following is an amino acid derived hormone ? | | | | |
|-----|------------|--|-----------------|--------------|--------------|---|---|---|---|-----|---|--|
| | belo | w: | | | | | | (1) | Estradiol | | | |
| | | Colur | | | | Column II | | (2) | Epinephrine | | | |
| | | (Fund | ction) | | | (Part of Excretory System) | tory | | (3) Estriol | | | |
| | a. | a. Ultrafiltration i. Henle' | | Henle's loop | | (4) | (4) Ecdysone | | | | | |
| | b. | Concentration of urine | | | ii. | Ureter | 49. | Which of the following hormones can play a | | | | |
| | c. | Transport of | | | iii. | Urinary bladder | | - | significant role in osteoporosis ?(1) Estrogen and Parathyroid hormone | | | |
| | | urine |) | | | | | (1) (2) | - | - | | |
| | d. | Stora | ige of uri | ne | iv. | Malpighian corpuscle | | (2) | | | | |
| | | | | | | - | | (4) | | | | |
| | | | | | v. | Proximal convoluted tubule | | (1) | Progesterone and Aldosterone | | | |
| | | a | b | c d | I | 50. | Which of the following structures or regions is | | | | | |
| | (1) | | | i | ii | | | inco | <i>prrectly</i> paired with | its | function ? | |
| | (1) (2) | v : | iv | ii | ii | | | (1) | Hypothalamus | : | production of | |
| | | iv | v | | ii | | | | | | releasing hormones and regulation of | |
| | (3) (4) | v iv | iv i | i ii | ii | | | | | | temperature, hunger and thirst. | |
| 47. | | atch the items given in Column I with those in lumn II and select the <i>correct</i> option given | | | | | | (2) | Medulla oblongata | : | controls respiration and cardiovascular reflexes. | |
| | 2010 | Colur | mn I | | Col | lumn II | | (3) | Corpus callosum | : | band of fibers connecting left and | |
| | a. | Glyco | osuria | i. | | umulation of uric in joints | | | | | right cerebral hemispheres. | |
| | b. | Gout | | ii. | | s of crystallised s within the kidney | | (4) | Limbic system | : | consists of fibre tracts that interconnect | |
| | c. | Rena | l calculi | iii. | | ammation in neruli | | | | | different regions of brain; controls | |
| | d. | Glom neph: | erular ritis | iv. | Pres urin | sence of glucose in e | | | | | movement. | |
| | | a | b | с | d | l | 51. | | transparent lens in place by | the | human eye is held in | |
| | (1) | ii | iii | i | iv | V | | (1) | smooth muscles att | ach | ed to the iris | |
| | (2) | iii | ii | iv | i | | | (1) shooth muscles attached to the fills (2) ligaments attached to the ciliary body (3) smooth muscles attached to the ciliary body | | | | |
| | (3) | iv | i | ii | ii | i | | | | | | |
| | (4) | i | ii | iii | iv | V | | (4) | ligaments attached | | | |

Downloade@4fepre.com

| 52. | stra | GTATCGCAT is a sequence from the coding and of a gene. What will be the corresponding sence of the transcribed mRNA? ACCUAUGCGAU AGGUAUCGCAU UCCAUAGCGUA UGGTUTCGCAT | | : |
|-----|--|---|---|----------|
| 53. | | of the following are part of an operon <i>except</i> an enhancer an operator a promoter structural genes | of many vertebrates is an example of (1) Convergent evolution (2) Homology (3) Adaptive radiation (4) Analogy 59. Which of the following is <i>not</i> an autoimmum | |
| 54. | Х | oman has an X-linked condition on one of her chromosomes. This chromosome can be erited by Only grandchildren Only daughters Both sons and daughters | disease ? (1) Alzheimer's disease (2) Psoriasis (3) Vitiligo (4) Rheumatoid arthritis 60. Which of the following characteristics representations of the following characteristics representation of the following c | |
| 55. | | Only sons ch the items given in Column I with those in iumn II and select the <i>correct</i> option given ow: <i>Column I</i> Proliferative Phase i. Breakdown of endometrial lining | c. Multiple allele d. Incomplete dominance e. Polygenic inheritance (1) b, d and e (2) b, c and e | |
| | b. c. (1) (2) (3) (4) | Secretory Phase ii. Follicular Phase Menstruation b c iii iii i iii ii i iii ii i iii iii | (3) a, c and e (4) a, b and c 61. Conversion of milk to curd improves it nutritional value by increasing the amount of (1) Vitamin B₁₂ (2) Vitamin D (3) Vitamin E (4) Vitamin A | ts |
| 56. | | i iii ii ording to Hugo de Vries, the mechanism of lution is Phenotypic variations Multiple step mutations Minor mutations Saltation | 62. In which disease does mosquito transmitter pathogen cause chronic inflammation of lymphatic vessels ? (1) Ringworm disease (2) Elephantiasis (3) Amoebiasis (4) Ascariasis | ed of |

Downloadest4Fernwover.com

| 63. | | | | | are inc | luded | in | 'Ex-situ | 68. | The | contracept |
|-----|---------|--|-------------------------|---------|------------------|---------|-------|----------------------|-----|------|-----------------------|
| | (1) | | tion' exce | - | | | | | | (1) | is an IUD |
| | (1) (2) | | anical ga llife safa | | l z a | | | | | (2) | blocks es |
| | (2) | | d banks | an par | K9 | | | | | | preventin |
| | (4) | | red grov | es | | | | | | (3) | is a post-o |
| | | | U | | | _ | | _ | | (4) | increases |
| 64. | | umn 🛛 | | - | | | | those in on given | | | prevents |
| | | Colı | ımn I | | (| Colum | n II | | 69. | The | amnion o |
| | a. | Eut | rophicat | ion | i. U | V-B ra | diat | tion | | fron | ı |
| | b. | San | itary lar | ndfill | ii. De | eforest | tatio | n | | (1) | mesodern |
| | c. | Sno | w blindr | ness | iii. N | utrien | t | | | (2) | ectoderm |
| | | | | | en | richm | ent | | | (3) | ectoderm |
| | d. | Jhu | m cultiv | ation | iv. W | aste d | ispo | sal | | (4) | endodern |
| | | a | b | С | d | | | | | (-) | 0114040111 |
| | (1) | iii | iv | i | ii | | | | 70. | The | differenc |
| | (2) | ii | i | iii | iv | | | | ••• | | miation is |
| | (3) | i | ii | iv | iii | | | | | - | |
| | (4) | i | iii | iv | ii | | | | | (1) | In sperm cells are |
| 65. | In a | a grow | ving pop | ulatior | n of a co | untry, | , | | | | seminifer |
| | (1) | | | | | | | | | | spermato |
| | (2) | | reprodu reprodu | | | | e m | ore than | | (2) | In sperm while in |
| | (3) | pre-reproductive individuals are less than the reproductive individuals. | | | | | | | | (3) | formed. In sperm |
| | (4) | - | oductive -reprodu | | | | ess t | than the | | | while in released |
| 66. | Whi | ch pa | rt of po | ppy pl | ant is 1 | used t | o ob | otain the | | | seminifer |
| | | | ack" ? | | | | | | | (4) | In sperm |

- (1)Roots
- (2)Flowers
- (3)Leaves
- (4)Latex
- 67. Which one of the following population interactions is widely used in medical science for the production of antibiotics ?
 - (1)Parasitism
 - (2)Commensalism
 - Amensalism (3)
 - Mutualism (4)

- tive 'SAHELI'
 - D.
- estrogen receptors in the uterus, ng eggs from getting implanted.
- -coital contraceptive.
- s the concentration of estrogen and ovulation in females.
- of mammalian embryo is derived
 - m and trophoblast
 - n and mesoderm
 - n and endoderm
 - m and mesoderm
- between spermiogenesis and ce 3
 - niogenesis spermatozoa from sertoli e released into the cavity of rous tubules, while in spermiation ozoa are formed.
 - niogenesis spermatids are formed, n spermiation spermatozoa are
 - niogenesis spermatozoa are formed, n spermiation spermatozoa are from sertoli cells into the cavity of rous tubules.
 - In spermiogenesis spermatozoa are formed, (4)while in spermiation spermatids are formed.
- 71. Hormones secreted by the placenta to maintain pregnancy are
 - hCG, hPL, progestogens, estrogens (1)
 - (2)hCG, hPL, progestogens, prolactin
 - (3)hCG, progestogens, estrogens, glucocorticoids
 - hCG, hPL, estrogens, relaxin, oxytocin (4)

| 72. | Column II and select the <i>correct</i> option given below : | | | | | | 75. | Col | umn I | | - | Column I with those in e <i>correct</i> option given |
|-----|--|--|--------------------|----------------------------|--------------|---------------------------------------|---------|----------------|----------|-----------|-----------|---|
| | | Colur | mn I | | (| Column II | | belo | ow: | | | |
| | a. | Tricu | ispid va | lve i | | etween left atrium | | | Colu | ımn I | | Column II |
| | 1. | D: | | ; | | nd left ventricle | | a. | Fibr | inogen | i. | Osmotic balance |
| | b. | Bicus | spid val | ve | | etween right entricle and | | b. | Glob | oulin | ii. | Blood clotting |
| | | | | | \mathbf{p} | ulmonary artery | | c. | Albu | ımin | iii. | Defence mechanism |
| | c. | Semi | lunar v | alve | | etween right | | | а | b | С | |
| | | | | atrium and right ventricle | | | (1) | i | iii | ii | | |
| | | a | b | с | | | | (2) | iii | ii | i | |
| | (1) | i | ii | iii | | | | (3) | ii | iii | i | |
| | (2) | iii | i | ii | | | | (4) | i | ii | iii | |
| | (3) | ii | i | iii | | | - | **** | | 0 1 | 6 11 . | |
| | (4) | i | iii | ii | | | 76. | | | | | g is an occupational |
| 73. | 8 | | | | | | - | | y disor | der : | | |
| | Column II and select the <i>correct</i> option given below : | | | (1) | | ılism | | | | | | |
| | | Colur | mn I | | | Column II i. 2500 – 3000 mL | | (2) | | nracis | | |
| | a. | Tidal | volum | е | i. | | | (3) | Emp | hysem | a | |
| | b. | Inspi | ratory] | Reserve | e ii | ii. 1100 – 1200 mL | | (4) | Silic | osis | | |
| | | volun | | | | | 77. | Cale | cium | is ir | nportant | in skeletal muscle |
| | c. | Expii volun | ratory H ne | Reserve | e ii | i. 500 – 550 mL | | con | | on beca | | |
| | d. | Resid | lual vol | ume | iv | . 1000 – 1100 mL | | (1) | | | he myosii | n head from the actin |
| | | a | b | с | d | | | (\mathbf{a}) | | nent. | | |
| | (1) | i | iv | ii | iii | | | (2) | | | - | remove the masking of for myosin. |
| | (2) | iii | ii | i | iv | | | (3) | | | | tion of bonds between |
| | (3) | iv | iii | ii | i | | | (3) | - | | | oridges and the actin |
| | (4) | iii | i | iv | ii | | | | filan | • | | 0 |
| 74. | Whi | | | follow | | options correctly ns in asthma and | | (4) | | vates tl | he myosin | ATPase by binding to |
| | - | | a, resp | - | | is in astima and | | | it. | | | |
| | (1) | | ased | respir | v | surface; | 78. | Whi | ich of | the f | ollowing | gastric cells indirectly |
| | (9) | Inflammation of bronchioles) Inflammation of bronchioles; Decrease | | | | helj | o in er | ythrop | oiesis ? | | | |
| | (2) | | nmatio ratory s | | | moles; Decreased | | (1) | Gobl | let cells | 5 | |
| | (3) | Decre | eased | respi | ratory | | | (2) | Chie | ef cells | | |
| | (A) | Inflammation of bronchioles | | | | (3) | Pari | etal cel | lls | | | |
| | (4) | | ratory s | | | nemoles; mcreased | | (4) | Muc | ous cel | ls | |

| 79. | Whi hom | ch one of these animals is <i>not</i> a eotherm? | 85. | Niss (1) | l bodies are mainly composed of Nucleic acids and SER |
|-----|--------------|---|-----|----------------|---|
| | (1) | Camelus | | (2) | Proteins and lipids |
| | (2) | Macropus | | (3) | Free ribosomes and RER |
| | (3) | Psittacula | | (4) | DNA and RNA |
| | (4) | Chelone | 86. | | ch of the following events does <i>not</i> occur in gh endoplasmic reticulum ? |
| 80. | | tify the vertebrate group of animals | | (1) | Cleavage of signal peptide |
| | char syst | acterized by crop and gizzard in its digestive | | (2) | Protein folding |
| | (1) | Aves | | (3) | Phospholipid synthesis |
| | (1) (2) | Amphibia | | (4) | Protein glycosylation |
| | (3) | Osteichthyes | 87. | Whi | ch of these statements is <i>incorrect</i> ? |
| | (4) | Reptilia | | (1) | Glycolysis operates as long as it is supplied with NAD that can pick up hydrogen atoms. |
| 81. | | tes differ from all other protozoans in | | (2) | Enzymes of TCA cycle are present in |
| | (1) | using pseudopodia for capturing prey | | (\mathbf{n}) | mitochondrial matrix. |
| | (2) | using flagella for locomotion | | (3) | Oxidative phosphorylation takes place in outer mitochondrial membrane. |
| | (3) | having two types of nuclei | | (4) | Glycolysis occurs in cytosol. |
| | (4) | having a contractile vacuole for removing excess water | 88. | Wh; | ch of the following terms describe human |
| | | | 00. | | tition ? |
| 82. | | ch of the following animals does <i>not</i> undergo amorphosis ? | | (1) | Pleurodont, Monophyodont, Homodont |
| | (1) | Moth | | (2) | Thecodont, Diphyodont, Homodont |
| | (2) | Earthworm | | (3) | Pleurodont, Diphyodont, Heterodont |
| | (3) | Starfish | | (4) | Thecodont, Diphyodont, Heterodont |
| | (4) | Tunicate | 89. | Man | y ribosomes may associate with a single |
| 83. | | ch of the following organisms are known as f producers in the oceans ? | | sim | NA to form multiple copies of a polypeptide ultaneously. Such strings of ribosomes are ned as |
| | (1) | Cyanobacteria | | (1) | Plastidome |
| | (2) | Dinoflagellates | | (2) | Polysome |
| | (3) | Euglenoids | | (3) | Nucleosome |
| | (4) | Diatoms | | (4) | Polyhedral bodies |
| 84. | | 0 | 90. | Sele | ct the <i>incorrect</i> match : |
| | | ale cockroach from a female cockroach ? | | (1) | Submetacentric – L-shaped chromososmes |
| | (1) | Forewings with darker tegmina | | (\mathbf{O}) | chromosomes |
| | (2) | Presence of a boat shaped sternum on the | | (2) | Lampbrush – Diplotene bivalents chromosomes |
| | | 9 th abdominal segment | | (3) | Polytene – Oocytes of amphibians |
| | (3) | Presence of anal cerci | | | chromosomes |
| | (4) | Presence of caudal styles | | (4) | Allosomes – Sex chromosomes |

| 91. | Pne | umatophores occur in | 98. | Which of the following pairs is <i>wrongly</i> | | | | | |
|-----|------|---|------|--|--|--|--|--|--|
| | (1) | Carnivorous plants | | matched ? | | | | | |
| | (2) | Halophytes | | (1) XO type sex : Grasshopper | | | | | |
| | (3) | Submerged hydrophytes | | determination | | | | | |
| | (4) | Free-floating hydrophytes | | (2) Starch synthesis in pea : Multiple alleles | | | | | |
| 92. | Swe | et potato is a modified | | (3) T.H. Morgan : Linkage (4) ABO blood grouping : Co-dominance | | | | | |
| | (1) | Tap root | | | | | | | |
| | (2) | Stem | 99. | Select the <i>correct</i> statement : | | | | | |
| | (3) | Rhizome | | (1) Spliceosomes take part in translation. | | | | | |
| | (4) | Adventitious root | | (2) Franklin Stahl coined the term "linkage". | | | | | |
| 93. | Whi | ich of the following statements is <i>correct</i> ? | | (3) Transduction was discovered by S. Altman. (4) Punnett square was developed by a British | | | | | |
| | (1) | Horsetails are gymnosperms. | | scientist. | | | | | |
| | (2) | Ovules are not enclosed by ovary wall in gymnosperms. | 100. | The experimental proof for semiconservative replication of DNA was first shown in a | | | | | |
| | (3) | Stems are usually unbranched in both <i>Cycas</i> and <i>Cedrus</i> . | | Plant Fungus | | | | | |
| | (4) | Selaginella is heterosporous, while Salvinia | | (3) Virus | | | | | |
| | | is homosporous. | | (4) Bacterium | | | | | |
| 94. | Cas | parian strips occur in | 101. | Select the <i>correct</i> match : | | | | | |
| | (1) | Cortex | | (1) Matthew Meselson – Pisum sativum | | | | | |
| | (2) | Epidermis | | and F. Stahl | | | | | |
| | (3) | Endodermis | | (2) Alec Jeffreys – Streptococcus | | | | | |
| | (4) | Pericycle | | pneumoniae | | | | | |
| 95. | | ondary xylem and phloem in dicot stem are | | (3) Francois Jacob and – Lac operon Jacques Monod | | | | | |
| | - | duced by | | (4) Alfred Hershey and – TMV | | | | | |
| | (1) | Phellogen | | Martha Chase | | | | | |
| | (2) | Apical meristems | 102. | Offsets are produced by | | | | | |
| | (3) | Axillary meristems | | (1) Parthenocarpy | | | | | |
| | (4) | Vascular cambium | | (2) Meiotic divisions | | | | | |
| 96. | Sele | ect the <i>wrong</i> statement : | | (3) Parthenogenesis | | | | | |
| | (1) | Pseudopodia are locomotory and feeding structures in Sporozoans. | | (4) Mitotic divisionsWhich of the following flowers only once in its | | | | | |
| | (2) | Cell wall is present in members of Fungi and Plantae. | | life-time ? (1) Mango | | | | | |
| | (3) | Mitochondria are the powerhouse of the cell in all kingdoms except Monera. | | (2) Bamboo species(3) Papaya | | | | | |
| | (4) | Mushrooms belong to Basidiomycetes. | | (4) Jackfruit | | | | | |
| 97. | Plar | nts having little or no secondary growth are | 104. | Which of the following has proved helpful in preserving pollen as fossils ? | | | | | |
| | (1) | Conifers | | (1) Oil content | | | | | |
| | (2) | Grasses | | (2) Pollenkitt | | | | | |
| | (3) | Cycads | | (3) Sporopollenin | | | | | |
| | (4) | Deciduous angiosperms | | (4) Cellulosic intine | | | | | |

| | (2) | 5 th June | | (2) | Extension, Denaturation, Annealing | | |
|------|-------|---|------|---|--|--|--|
| | | | | (3) | Denaturation, Annealing, Extension | | |
| | (3) | 22 nd April | | (4) | Annealing, Extension, Denaturation | | |
| | (4) | 21 st April | 119 | | | | |
| 106. | Nata | ality refers to | 112. | | India, the organisation responsible for essing the safety of introducing genetically | | |
| | (1) | Number of individuals leaving the habitat | | | dified organisms for public use is | | |
| | (2) | Death rate | | (1) | Research Committee on Genetic | | |
| | (3) | Number of individuals entering a habitat | | | Manipulation (RCGM) | | |
| | (4) | Birth rate | | (2) | Indian Council of Medical Research (ICMR) | | |
| 107. | | t type of ecological pyramid would be ined with the following data? | | (3) | Genetic Engineering Appraisal Committee (GEAC) | | |
| | | Secondary consumer : 120 g | 113. | (4) | Council for Scientific and Industrial Research (CSIR) | | |
| | | Primary consumer : 60 g | | | | | |
| | | Primary producer : 10 g | | | ich of the following is commonly used as a | | |
| | (1) | Upright pyramid of numbers | | | tor for introducing a DNA fragment in human aphocytes ? | | |
| | (2) | Inverted pyramid of biomass | | (1) | λ phage | | |
| | (3) | Upright pyramid of biomass | | (2) | Retrovirus | | |
| | (4) | Pyramid of energy | | (2) | pBR 322 | | |
| 108. | Whi | ch of the following is a secondary pollutant? | | (4) | Ti plasmid | | |
| 1000 | (1) | SO_2 | | | - | | |
| | | - | 114. | | e of bioresources by multinational companies l organisations without authorisation from the | | |
| | (2) | СО | | concerned country and its people is called | | | |
| | (3) | O_3 | | (1) | Biodegradation | | |
| | (4) | CO_2 | | (2) | Bio-infringement | | |
| 109. | Nich | ie is | | (3) | Bioexploitation | | |
| | (1) | the range of temperature that the organism needs to live | | (4) | Biopiracy | | |
| | (2) | all the biological factors in the organism's environment | 115. | A 'new' variety of rice was patented by a foreign company, though such varieties have been present in India for a long time. This is related to | | | |
| | (3) | the functional role played by the organism where it lives | | (1) | Lerma Rojo | | |
| | (4) | the physical space where an organism lives | | (2) | Co-667 | | |
| 110. | In st | tratosphere, which of the following elements | | (3) (4) | Basmati Sharbati Sonora | | |
| | acts | as a catalyst in degradation of ozone and | | | | | |
| | | ase of molecular oxygen ? | 116. | | ect the <i>correct</i> match : | | |
| | (1) | Fe | | (1) | T.H. Morgan – Transduction | | |
| | (2) | Carbon | | (2) | Ribozyme – Nucleic acid | | |
| | (3) | Oxygen | | (3) | G. Mendel – Transformation | | |

111. The correct order of steps in Polymerase Chain

Denaturation, Extension, Annealing

Reaction (PCR) is

(1)

105. World Ozone Day is celebrated on

16th September

(1)

(4)

Cl

(4)

 $F_2 \times Recessive parent$

English

Dihybrid cross

—

| 115 | | 104 | |
|------|--|------|---|
| 117. | Which of the following elements is responsible for maintaining turgor in cells ? | 124. | |
| | (1) Potassium | | (1) Respiration in bacteria |
| | (1) Fotassium (2) Magnesium | | (2) Fatty acid breakdown |
| | (3) Calcium | | (3) Activation of amino acid |
| | (4) Sodium | | (4) Formation of secretory vesicles |
| 110 | | 125. | The two functional groups characteristic of |
| 118. | Which one of the following plants shows a very close relationship with a species of moth, where | | sugars are |
| | none of the two can complete its life cycle without | | (1) carbonyl and phosphate |
| | the other ? | | (2) hydroxyl and methyl |
| | (1) Banana | | (3) carbonyl and hydroxyl |
| | (2) Hydrilla | | (4) carbonyl and methyl |
| | (3) Viola | 126. | Which among the following is <i>not</i> a prokaryote ? |
| | (4) Yucca | | (1) Nostoc |
| 119. | Pollen grains can be stored for several years in | | (2) Saccharomyces |
| - | liquid nitrogen having a temperature of | | (3) Oscillatoria |
| | (1) $-196^{\circ}C$ | | (4) Mycobacterium |
| | (2) $-120^{\circ}C$ | 127. | Stomatal movement is <i>not</i> affected by |
| | (3) $-160^{\circ}C$ | | (1) O_2 concentration |
| | (4) $-80^{\circ}C$ | | (2) Temperature |
| 120 | Double fertilization is | | (3) CO_2 concentration |
| 120. | (1) Fusion of two male gametes with one egg | | (4) Light |
| | (1) Fusion of two male gametes of a pollen tube | 128. | Which of the following is not a product of light |
| | with two different eggs | | reaction of photosynthesis ? |
| | (3) Syngamy and triple fusion | | (1) NADPH |
| | (4) Fusion of one male gamete with two polar | | (2) ATP |
| | nuclei | | (3) Oxygen |
| 121. | Oxygen is ${\it not}$ produced during photosynthesis by | | (4) NADH |
| | (1) Cycas | 129. | Which of the following is true for nucleolus ? |
| | (2) Green sulphur bacteria | | (1) It takes part in spindle formation. |
| | (3) Chara | | (2) Larger nucleoli are present in dividing cells. |
| | (4) Nostoc | | (3) It is a site for active ribosomal RNA |
| 122. | What is the role of NAD ⁺ in cellular | | synthesis. |
| | respiration ? | | (4) It is a membrane-bound structure. |
| | (1) It is a nucleotide source for ATP synthesis. | 130. | The stage during which separation of the paired |
| | (2) It functions as an enzyme. | | homologous chromosomes begins is |
| | (3) It is the final electron acceptor for anaerobic | | (1) Diakinesis |
| | respiration. | | (2) Pachytene |
| | (4) It functions as an electron carrier. | | (3) Zygotene |
| 123. | In which of the following forms is iron absorbed | | (4) Diplotene |
| | by plants ? | 131. | 5 |
| | (1) Free element | | (1) Rectangular |
| | (2) Ferric | | (2) Dumb-bell shaped |
| | (3) Both ferric and ferrous | | (3) Barrel shaped |
| | (4) Ferrous | | (4) Kidney shaped |

CHLAA/HH/Page 15

Downloadest/feernewswewswekshapter.com

| 132. | | mn II | | - | n in Column I with those in t the <i>correct</i> option given | | | | | |
|-------|---------|--------------------|--------|--------|--|------|--------|--------|---------|--|
| | | Colum | n I | | Colun | nn l | Ί | | | |
| | a. | Herba | rium | i. | It is a collect plants | ved | | | | |
| | b. | Key | | ii. | A list that enumerates methodically all the species found in an area with brief description aiding identification. | | | | | |
| | c. | Museu | ım | iii. | pressed plant specimens mounted on sheets are kept. | | | | | |
| | d. | Catalo | gue | iv. | | | | | | |
| | | a | b | с | d | _ | | | | |
| | (1) | ii | iv | iii | i | - | | | | |
| | (2) | i | iv | iii | ii | | | | | |
| | (3) | iii | iv | i | ii | - | | | | |
| | (4) | iii | ii | i | iv | | | | | |
| | | | | | | | | | | |
| 133. | | ch one i | | | matche | ed ' | | 1 | | |
| | (1) | Gemm | - | | | _ | | chant | | |
| | (2) | Unifla | 0 | 0 | | _ | • | sipho | onia | |
| | (3) | Unicel | | | | _ | | orella | | |
| | (4) | Billage | enate | zoosp | ores - | _ | Brov | vn alş | gae | |
| 134. | | r karyo uced ex | | | | me | eiosis | s, spo | res are | |
| | (1) | Agaric | us | | | | | | | |
| | (2) | Neuros | spora | | | | | | | |
| | (3) | Sacche | aromy | ces | | | | | | |
| | (4) | Altern | aria | | | | | | | |
| 135. | Wing | ged poll | en gra | ains a | re pres | sent | tin | | | |
| 1.000 | (1) | Mango | - | | -0 proc | | | | | |
| | (1) (2) | - | | | | | | | | |
| | | Pinus | | | | | | | | |
| | (4) | Cycas | | | | | | | | |

136. In which case is the number of molecules of water maximum ?

- (1) 0.00224 L of water vapours at 1 atm and 273 K
- $(2) \quad 18 \ \text{mL of water} \\$
- (3) 10^{-3} mol of water
- $(4) \quad 0.18 \ g \ of \ water$
- **137.** The correct difference between first- and second-order reactions is that
 - (1) a first-order reaction can be catalyzed; a second-order reaction cannot be catalyzed
 - (2) the rate of a first-order reaction does not depend on reactant concentrations; the rate of a second-order reaction does depend on reactant concentrations
 - (3) the rate of a first-order reaction does depend on reactant concentrations; the rate of a second-order reaction does not depend on reactant concentrations
 - (4) the half-life of a first-order reaction does not depend on $[A]_0$; the half-life of a second-order reaction does depend on $[A]_0$
- **138.** Consider the change in oxidation state of Bromine corresponding to different emf values as shown in the diagram below :

Then the species undergoing disproportionation is

- (1) Br₂
- (2) BrO_{3}^{-}
- (3) HBrO
- (4) BrO₄⁻
- **139.** Among CaH_2 , BeH_2 , BaH_2 , the order of ionic character is
 - (1) $BeH_2 < BaH_2 < CaH_2$
 - (2) $\operatorname{BeH}_2 < \operatorname{CaH}_2 < \operatorname{BaH}_2$
 - $(3) \quad \operatorname{BaH}_2 < \operatorname{BeH}_2 < \operatorname{CaH}_2$
 - $(4) \quad \mathrm{CaH}_2 < \mathrm{BeH}_2 < \mathrm{BaH}_2$

| | (1) | MnO | • | | | | (1) | Any opecting have $1 \rightarrow 4$ d-mixage and $1 \rightarrow 6 \beta$ -linkage | | | | |
|------|--|--------------------------|-----------------------|------------|--|--------------|---|--|--|--|--|--|
| | (2) | CrO_4^2 | | | | | (2) | Amylopectin have $1 \rightarrow 4 \alpha$ -linkage and $1 \rightarrow 6 \alpha$ -linkage | | | | |
| | (3) | MnO | $\frac{2}{4}$ | | | | (3) | Amylose is made up of glucose and | | | | |
| | (4) | Cr_2O | $\frac{2}{7}$ | | | | | galactose | | | | |
| 141. | Iron | carbon | nyl, Fe(C | $(O)_5$ is | | | (4) | $\begin{array}{llllllllllllllllllllllllllllllllllll$ | | | | |
| | (1) | trinuc | clear | | | 146. | Reg | Regarding cross-linked or network polymers, | | | | |
| | (2) | tetrar | nuclear | | | 1100 | - | which of the following statements is <i>incorrect</i> ? | | | | |
| | (3) | dinuc | | | | | (1) | (1) Examples are bakelite and melamine. | | | | |
| | (4) | mononuclear | | | | | (2) | | | | | |
| 142. | | | | - | n in Column I with the | | | various linear polymer chains. | | | | |
| | _ | - | | | of the ions given in <i>correct</i> code : | | (3) | They contain strong covalent bonds in their polymer chains. | | | | |
| | | Colun | nn I | | Column II | | (4) | They are formed from bi- and tri-functional | | | | |
| | a. | Co ³⁺ | | i. | $\sqrt{8}$ B.M. | | | monomers. | | | | |
| | b. | Cr ³⁺ | | ii. | $\sqrt{35}$ B.M. | 147. | | ation of aniline in strong acidic medium also | | | | |
| | | Fe ³⁺ | | | | | give (1) | s m-nitroaniline because In absence of substituents nitro group | | | | |
| | c. | | | iii. | $\sqrt{3}$ B.M. | | (1) | always goes to m-position. | | | | |
| | d. | Ni^{2+} | | iv. | $\sqrt{24}$ B.M. | | (2) | In spite of substituents nitro group always | | | | |
| | | | | v. | $\sqrt{15}$ B.M. | | | goes to only m-position. | | | | |
| | | a | b | С | d | | (3) | In acidic (strong) medium aniline is present as anilinium ion. | | | | |
| | (1) | iv | i | ii | iii | | (4) | In electrophilic substitution reactions | | | | |
| | (2) | iv | v | ii | i | | | amino group is meta directive. | | | | |
| | (3) | iii | v | i | ii | 148. | Which of the following oxides is most acidic in nature? | | | | | |
| | (4) | i | ii | iii | iv | | (1) | BaO | | | | |
| 143. | | - | | - | netic behaviour of the | | (2) | MgO | | | | |
| | com | plex [N | [i(CO) ₄] | are | | | (3) | CaO | | | | |
| | (1) | _ | - | - | try and paramagnetic | | (4) | BeO | | | | |
| | (2) | _ | - | - | try and diamagnetic | 149. | Am | ixture of 2.3 g formic acid and 4.5 g oxalic | | | | |
| | (3) | | - | | and paramagnetic | 140. | | is treated with conc. H_2SO_4 . The evolved | | | | |
| | (4) | | - | | v and diamagnetic | | | eous mixture is passed through KOH pellets. | | | | |
| 144. | The type of isomerism shown by the complex $[CoCl_2(en)_2]$ is | | | | | | Weig will | ght (in g) of the remaining product at STP be | | | | |
| | (1) Ionization isomerism | | | | | | (1) | 2.8 | | | | |
| | (2) | Geom | etrical i | someris | sm | | (2) | 1.4 | | | | |
| | (3) | Linka | ige isom | erism | | | (3) | 4.4 | | | | |
| | (4) | Coord | lination | isomeri | sm | | (4) | 3.0 | | | | |
| CHL | AA/HF | l/Page | 17 | Dov | vnloade@1460FARv | <u>80101</u> | 4 <mark>00</mark> 4 | whapter.com English | | | | |

140. Which one of the following ions exhibits 145. The difference between amylose and amylopectin

is (1)

Amylopectin have 1 \rightarrow 4 $\alpha\text{-linkage}$ and

d-d transition and paramagnetism as well?

(1)

 MnO_4^-


- 150. Hydrocarbon (A) reacts with bromine by substitution to form an alkyl bromide which by Wurtz reaction is converted to gaseous hydrocarbon containing less than four carbon atoms. (A) is
 - (1) $CH_3 CH_3$
 - (2) $CH \equiv CH$
 - (3) CH₄
 - (4) $CH_2 = CH_2$
- **151.** Which oxide of nitrogen is *not* a common pollutant introduced into the atmosphere both due to natural and human activity ?
 - (1) N₂O
 - $(2) N_2O_5$
 - (3) NO
 - (4) NO₂
- 152. The compound C_7H_8 undergoes the following reactions :

$$\mathrm{C_7H_8} \xrightarrow{3 \mathrm{Cl}_2 / \Delta} \mathrm{A} \xrightarrow{\mathrm{Br}_2 / \mathrm{Fe}} \mathrm{B} \xrightarrow{\mathrm{Zn} / \mathrm{HCl}} \mathrm{C}$$

The product 'C' is

- $(1) \quad \ \ 3\text{-}bromo-2,4,6\text{-}trichlorotoluene}$
- (2) *m*-bromotoluene
- (3) *p*-bromotoluene
- (4) o-bromotoluene
- 153. The compound A on treatment with Na gives B, and with PCl₅ gives C. B and C react together to give diethyl ether. A, B and C are in the order
 - (1) $C_2H_5Cl, C_2H_6, C_2H_5OH$
 - $(2) \quad C_2H_5OH, C_2H_6, C_2H_5Cl$
 - (3) C_2H_5OH , C_2H_5ONa , C_2H_5Cl
 - (4) C_2H_5OH , C_2H_5Cl , C_2H_5ONa

by **154.** Which of the following carbocations is expected to be most stable ?



- **155.** Which of the following is correct with respect to -I effect of the substituents ? (R = alkyl)
 - (1) $-NH_2 > -OR > -F$
 - (2) $-NH_2 < -OR < -F$
 - (3) $-NR_2 > -OR > -F$
 - $(4) \quad -\mathrm{NR}_2 < -\mathrm{OR} < -\mathrm{F}$
- **156.** Which of the following molecules represents the order of hybridisation sp², sp², sp, sp from left to right atoms ?
 - (1) $CH_2 = CH CH = CH_2$
 - (2) $HC \equiv C C \equiv CH$
 - $(3) \quad \mathrm{CH}_3 \mathrm{CH} = \mathrm{CH} \mathrm{CH}_3$
 - (4) $CH_2 = CH C \equiv CH$

- 157. Carboxylic acids have higher boiling points than 160. Identify the major products P, Q and R in the and alcohols aldehydes, ketones even of comparable molecular mass. It is due to their
 - (1)more extensive association of carboxylic acid via van der Waals force of attraction
 - (2)formation of intramolecular H-bonding
 - (3)formation of intermolecular H-bonding
 - (4)formation of carboxylate ion
- 158. Compound A, $C_8H_{10}O$, is found to react with NaOI (produced by reacting Y with NaOH) and yields a yellow precipitate with characteristic smell.

A and Y are respectively

(1)
$$CH - CH_3 \text{ and } I_2$$


 $I \\ OH$

(2)
$$H_3C \longrightarrow CH_2 - OH \text{ and } I_2$$

(3)
$$CH_3 \longrightarrow OH \text{ and } I_2$$

(4)
$$\bigcirc$$
 CH₂ – CH₂ – OH and I₂

159. In the reaction



the electrophile involved is

- Θ dichloromethyl anion (CHCl₂) (1)
- (2)dichloromethyl cation $(CHCl_2)$
- (3)dichlorocarbene (:CCl₂)
- \oplus formyl cation (CHO) (4)

following sequence of reactions :

$$\begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} Anhydrous \\ AICl_{3} \end{array} \end{array} \\ P \\ \begin{array}{c} \begin{array}{c} (i) \\ O_{2} \\ (ii) \\ H_{3}O^{+/\Delta} \end{array} Q + R \end{array} \end{array} \\ \begin{array}{c} P \\ \begin{array}{c} P \\ Q \\ R \end{array} \end{array} \\ \begin{array}{c} \begin{array}{c} P \\ (1) \end{array} \\ \begin{array}{c} \begin{array}{c} \begin{array}{c} CH(CH_{3})_{2} \\ , \end{array} \end{array} \\ \begin{array}{c} \begin{array}{c} OH \\ , \end{array} \end{array} \\ \begin{array}{c} CH_{2}CH_{2}CH_{3} \\ , \end{array} \\ \begin{array}{c} \begin{array}{c} CH_{2}CH_{2}CH_{3} \\ , \end{array} \\ \begin{array}{c} \begin{array}{c} CH_{3} \\ , \end{array} \end{array} \\ \begin{array}{c} \begin{array}{c} CH_{2}CH_{2}CH_{3} \\ , \end{array} \\ \begin{array}{c} \begin{array}{c} CH_{3} \\ , \end{array} \\ \begin{array}{c} CH_{3}CH_{2} - OH \\ , \end{array} \\ \begin{array}{c} CH_{3}CH_{2} - OH \\ \end{array} \\ \begin{array}{c} \begin{array}{c} CH_{3}CH_{3} - CO - CH_{3} \end{array} \\ \end{array} \\ \begin{array}{c} \begin{array}{c} \end{array} \\ \begin{array}{c} \begin{array}{c} CH_{2}CH_{2}CH_{3} \\ , \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \begin{array}{c} CH_{2}CH_{2}CH_{3} \\ , \end{array} \\ \begin{array}{c} CH_{3} - CO - CH_{3} \end{array} \end{array} \end{array}$$
 \\ \begin{array}{c} \begin{array}{c} CH_{3} \\ \end{array} \\ \begin{array}{c} CH_{2}CH_{2}CH_{3} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \begin{array}{c} CH_{2}CH_{2}CH_{3} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \end{array} \\ \begin{array}{c} CH_{2} \\ \end{array} \\ \begin{array}{c} CH_{2}CH_{2}CH_{3} \\ \end{array} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \end{array} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ \end{array} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ CH_{3} \\ \\ \end{array} \\ \end{array} \\ \end{array} \\ \begin{array}{c} CH_{3} \\ CH_{3} \\ \end{array} \\ \\ \end{array} \\ \begin{array}{c} CH_{3} \\ CH_{3} \\

- 161. Which of the following compounds can form a zwitterion?
 - (1)Benzoic acid
 - Aniline (2)
 - (3)Glycine
 - (4)Acetanilide

162. The correct order of atomic radii in group 13 | 168. For the redox reaction elements is

- (1) B < Ga < Al < Tl < In
- (2)B < Al < In < Ga < Tl
- B < Ga < Al < In < Tl(3)
- B < Al < Ga < In < Tl(4)
- 163. Which one of the following elements is unable to form MF_6^{3-} ion ?
 - В (1)
 - (2)Ga
 - (3)In
 - (4)Al
- 164. The correct order of N-compounds in its decreasing order of oxidation states is
 - HNO₃, NH₄Cl, NO, N₂ (1)
 - (2)HNO₃, NO, N₂, NH₄Cl
 - NH₄Cl, N₂, NO, HNO₃ (3)
 - HNO₃, NO, NH₄Cl, N₂ (4)
- **165.** Which of the following statements is *not* true for halogens?
 - (1)All but fluorine show positive oxidation states.
 - (2)All form monobasic oxyacids.
 - (3)Chlorine has the highest electron-gain enthalpy.
 - (4)All are oxidizing agents.
- **166.** In the structure of ClF_3 , the number of lone pairs of electrons on central atom 'Cl' is
 - (1)four
 - (2)one
 - (3)three
 - (4)two
- 167. Considering Ellingham diagram, which of the following metals can be used to reduce alumina?
 - (1)Mg
 - Fe (2)
 - (3)Cu
 - (4)Zn

 $MnO_4^- + C_2O_4^{2-} + H^+ \longrightarrow Mn^{2+} + CO_2 + H_2O_4^$ the correct coefficients of the reactants for the balanced equation are

| | MnO_4^- | $C_2 O_4^{2-}$ | H^{+} |
|-----|--------------------|----------------|------------------|
| (1) | 2 | 16 | 5 |
| (2) | 16 | 5 | 2 |
| (3) | 5 | 16 | 2 |
| (4) | 2 | 5 | 16 |

- 169. The bond dissociation energies of X_2 , Y_2 and XYare in the ratio of 1: 0.5: 1. ΔH for the formation of XY is -200 kJ mol^{-1} . The bond dissociation energy of X₂ will be
 - 800 kJ mol^{-1} (1)
 - 200 kJ mol^{-1} (2)
 - 400 kJ mol^{-1} (3)
 - 100 kJ mol^{-1} (4)
- 170. The correction factor 'a' to the ideal gas equation corresponds to
 - (1)electric field present between the gas molecules
 - (2)density of the gas molecules
 - forces of attraction between the (3)gas molecules
 - (4)volume of the gas molecules
- 171. Which one of the following conditions will favour maximum formation of the product in the reaction.

 $A_2(g) + B_2(g) \rightleftharpoons X_2(g) \quad \Delta_r H = -X kJ?$

- High temperature and high pressure (1)
- (2)Low temperature and high pressure
- (3)High temperature and low pressure
- (4)Low temperature and low pressure
- **172.** When initial concentration of the reactant is doubled, the half-life period of a zero order reaction
 - (1)is tripled
 - (2)is halved
 - remains unchanged (3)
 - (4)is doubled

173. Given van der Waals constant for NH_3 , H_2 , O_2 and CO_2 are respectively 4.17, 0.244, 1.36 and 3.59, which one of the following gases is most easily liquefied ?

- (1) O_2
- (2) NH_3
- CO_{2} (3)
- H_{2} (4)
- 174. Following solutions were prepared by mixing 178. Iron exhibits bcc structure at room temperature. different volumes of NaOH and HCl of different concentrations :
 - 60 mL $\frac{M}{10}$ HCl + 40 mL $\frac{M}{10}$ NaOH a. 55 mL $\frac{M}{10}$ HCl + 45 mL $\frac{M}{10}$ NaOH b. 75 mL $\frac{M}{5}$ HCl + 25 mL $\frac{M}{5}$ NaOH c. $100 \text{ mL} \frac{\text{M}}{10} \text{ HCl} + 100 \text{ mL} \frac{\text{M}}{10} \text{ NaOH}$ d. pH of which one of them will be equal to 1? (1)d

 - (2)b
 - (3)с
 - (4)а
- 175. On which of the following properties does the coagulating power of an ion depend?
 - (1)Both magnitude and sign of the charge on the ion
 - (2)The magnitude of the charge on the ion alone
 - (3)The sign of charge on the ion alone
 - (4)Size of the ion alone

176. The solubility of $BaSO_4$ in water is $2{\cdot}42\times10^{-3}~{\rm gL}^{-1}$ at 298 K. The value of its solubility product (K_{sp}) will be

(Given molar mass of $BaSO_4 = 233 \text{ g mol}^{-1}$)

- $1.08 \times 10^{-14} \text{ mol}^2 \text{ L}^{-2}$ (1)
- $1.08 imes 10^{-10} \text{ mol}^2 \text{ L}^{-2}$ (2)
- $1.08 imes 10^{-8} ext{ mol}^2 ext{ L}^{-2}$ (3)
- $1.08 \times 10^{-12} \text{ mol}^2 \text{ L}^{-2}$ (4)

- Magnesium reacts with an element (X) to form an 177. ionic compound. If the ground state electronic configuration of (X) is $1s^2 2s^2 2p^3$, the simplest formula for this compound is
 - (1)Mg₂X
 - (2) $Mg_{2}X_{3}$
 - Mg₂X₂ (3)
 - MgX₂ (4)
 - Above 900°C, it transforms to fcc structure. The ratio of density of iron at room temperature to that at 900°C (assuming molar mass and atomic radii of iron remains constant with temperature) is

(1)
$$\frac{3\sqrt{3}}{4\sqrt{2}}$$

(2)
$$\frac{\sqrt{3}}{\sqrt{2}}$$

(3)
$$\frac{1}{2}$$

(4)
$$\frac{4\sqrt{3}}{3\sqrt{2}}$$

179. Which one is a *wrong* statement?

The electronic configuration of N atom is (1)

$$\begin{array}{c} 1s^2 \\ \uparrow \downarrow \end{array} \begin{array}{c} 2s^2 \\ \uparrow \downarrow \end{array} \begin{array}{c} 2p_x^1 \ 2p_y^1 \ 2p_z^1 \\ \uparrow \downarrow \end{array} \end{array}$$

- (2)Total orbital angular momentum of electron in 's' orbital is equal to zero.
- (3)The value of m for d_{z2} is zero.
- (4)An orbital is designated by three quantum numbers while an electron in an atom is designated by four quantum numbers.
- **180.** Consider the following species :

CN⁺, CN⁻, NO and CN

Which one of these will have the highest bond order?

- (1) CN^+
- (2)NO
- (3)CN

SPACE FOR ROUGH WORK

SPACE FOR ROUGH WORK

Read carefully the following instructions :

- 1. Each candidate must show on demand his/her Admit Card to the Invigilator.
- 2. No candidate, without special permission of the Superintendent or Invigilator, would leave his/her seat.
- 3. The candidates should not leave the Examination Hall without handing over their Answer Sheet to the Invigilator on duty and sign the Attendance Sheet twice. **Cases where a** candidate has not signed the Attendance Sheet second time will be deemed not to have handed over the Answer Sheet and dealt with as an unfair means case.
- 4. Use of Electronic/Manual Calculator is prohibited.
- 5. The candidates are governed by all Rules and Regulations of the examination with regard to their conduct in the Examination Hall. All cases of unfair means will be dealt with as per Rules and Regulations of this examination.
- 6. No part of the Test Booklet and Answer Sheet shall be detached under any circumstances.
- 7. The candidates will write the Correct Test Booklet Code as given in the Test Booklet/Answer Sheet in the Attendance Sheet.