HLAAC

Test Booklet Code

This Booklet contains 24 pages.



Do not open this Test Booklet until you are asked to do so.

Read carefully the Instructions on the Back Cover of this Test Booklet.


Important Instructions :

- 1. The Answer Sheet is inside this Test Booklet. When you are directed to open the Test Booklet, take out the Answer Sheet and fill in the particulars on **Side-1** and **Side-2** carefully with **blue/black** ball point pen only.
- 2. The test is of **3 hours** duration and this Test Booklet contains **180** questions. Each question carries **4** marks. For each correct response, the candidate will get **4** marks. For each incorrect response, **one mark** will be deducted from the total scores. The maximum marks are 720.
- 3. Use Blue/Black Ball Point Pen only for writing particulars on this page/marking responses.
- 4. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
- 5. On completion of the test, the candidate must hand over the Answer Sheet to the Invigilator before leaving the Room/Hall. The candidates are allowed to take away this Test Booklet with them.
- 6. The CODE for this Booklet is **KK**. Make sure that the CODE printed on **Side-2** of the Answer Sheet is the same as that on this Test Booklet. In case of discrepancy, the candidate should immediately report the matter to the Invigilator for replacement of both the Test Booklet and the Answer Sheet.
- 7. The candidates should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your Roll No. anywhere else except in the specified space in the Test Booklet/Answer Sheet.
- 8. Use of white fluid for correction is *not* permissible on the Answer Sheet.

Name of the Candidate (in Capitals) :	
Roll Number : in figures	
: in words	
Centre of Examination (in Capitals) :	
Candidate's Signature :	Invigilator's Signature :
Facsimile signature stamp of Centre Superintendent :	

HLAAC/KK/Page 1 Downloaded from www.cclchapter.com

1. The kinetic energies of a planet in an elliptical orbit about the Sun, at positions A, B and C are K_A , K_B and K_C , respectively. AC is the major axis and SB is perpendicular to AC at the position of the Sun S as shown in the figure. Then




- (1) $K_A > K_B > K_C$
- (2) $K_B < K_A < K_C$
- (3) $K_A < K_B < K_C$
- $(4) \quad \mathbf{K}_{\mathbf{B}} > \mathbf{K}_{\mathbf{A}} > \mathbf{K}_{\mathbf{C}}$
- - (1) 5:7
 - (2) 10:7
 - (3) 7:10
 - (4) 2:5
- **3.** A solid sphere is rotating freely about its symmetry axis in free space. The radius of the sphere is increased keeping its mass same. Which of the following physical quantities would remain constant for the sphere ?
 - (1) Moment of inertia
 - (2) Rotational kinetic energy
 - (3) Angular velocity
 - (4) Angular momentum
- 4. If the mass of the Sun were ten times smaller and the universal gravitational constant were ten times larger in magnitude, which of the following is *not* correct ?
 - (1) Walking on the ground would become more difficult.
 - (2) Time period of a simple pendulum on the Earth would decrease.
 - (3) Raindrops will fall faster.
 - (4) 'g' on the Earth will not change.

- A toy car with charge q moves on a frictionless horizontal plane surface under the influence of a uniform electric field \vec{E} . Due to the force q \vec{E} , its velocity increases from 0 to 6 m/s in one second duration. At that instant the direction of the field is reversed. The car continues to move for two more seconds under the influence of this field. The average velocity and the average speed of the toy car between 0 to 3 seconds are respectively
 - (1) 1 m/s, 3 m/s
 - (2) 1 m/s, 3.5 m/s
 - (3) 2 m/s, 4 m/s

6.

7.

- (4) 1.5 m/s, 3 m/s
- A block of mass m is placed on a smooth inclined wedge ABC of inclination θ as shown in the figure. The wedge is given an acceleration 'a' towards the right. The relation between a and θ for the block to remain stationary on the wedge is



(2)
$$a = g \cos \theta$$

(3)
$$a = \frac{g}{\operatorname{cosec} \theta}$$

(4) $a = g \tan \theta$


The moment of the force, $\vec{F} = 4\hat{i} + 5\hat{j} - 6\hat{k}$ at (2, 0, -3), about the point (2, -2, -2), is given by

(1) $-4\hat{i} - \hat{j} - 8\hat{k}$ (2) $-7\hat{i} - 8\hat{j} - 4\hat{k}$ (3) $-8\hat{i} - 4\hat{j} - 7\hat{k}$

$$(4) \quad -7\, {\hat i} \ -4\, {\hat j} \ -8\, {\hat k}$$

- A student measured the diameter of a small steel ball using a screw gauge of least count 0.001 cm. The main scale reading is 5 mm and zero of circular scale division coincides with 25 divisions above the reference level. If screw gauge has a zero error of -0.004 cm, the correct diameter of the ball is
 - (1) 0.525 cm(2) 0.053 cm
 - (3) 0.521 cm
 - (4) 0.521 cm

9. The volume (V) of a monatomic gas varies with its temperature (T), as shown in the graph. The ratio of work done by the gas, to the heat absorbed by it, when it undergoes a change from state A to state B, is



- 10. The fundamental frequency in an open organ pipe is equal to the third harmonic of a closed organ pipe. If the length of the closed organ pipe is 20 cm, the length of the open organ pipe is
 - (1) 8 cm
 - $(2) \quad 12.5 \text{ cm}$
 - $(3) \quad 13{\cdot}2\ cm$
 - (4) 16 cm
- **11.** At what temperature will the rms speed of oxygen molecules become just sufficient for escaping from the Earth's atmosphere ? (Given :


Mass of oxygen molecule (m) = 2.76×10^{-26} kg Boltzmann's constant k_B = 1.38×10^{-23} J K⁻¹)


- $(1) \quad 8{\cdot}360 \times 10^4 \; \mathrm{K}$
- (2) $5.016 \times 10^4 \text{ K}$
- (3) $2.508 \times 10^4 \text{ K}$
- $(4) \quad 1{\cdot}254\times 10^4 \ \mathrm{K}$
- 12. The efficiency of an ideal heat engine working between the freezing point and boiling point of water, is
 - (1) 20%
 - $(2) \quad 6.25\%$
 - (3) 26.8%
 - (4) 12.5%

- A carbon resistor of (47 ± 4.7) k Ω is to be marked with rings of different colours for its identification. The colour code sequence will be
 - $(1) \quad Yellow-\ Violet-Orange-Silver$
 - $(2) \quad Yellow-\ Green-Violet-Gold$
 - (3) Violet Yellow Orange Silver
 - (4) Green Orange Violet Gold
- 14. A set of 'n' equal resistors, of value 'R' each, are connected in series to a battery of emf 'E' and internal resistance 'R'. The current drawn is I. Now, the 'n' resistors are connected in parallel to the same battery. Then the current drawn from battery becomes 10 I. The value of 'n' is
 - (1) 11
 - (2) 20
 - (3) 10
 - (4) 9
- 15. A battery consists of a variable number 'n' of identical cells (having internal resistance 'r' each) which are connected in series. The terminals of the battery are short-circuited and the current I is measured. Which of the graphs shows the correct relationship between I and n?

(1)
$$\uparrow$$
 \longrightarrow n

τl





Downloadeet4fernweverwetehapter.com

- 16. An em wave is propagating in a medium with a velocity \$\vec{V}\$ = V\$\$\$\$\$\$\$\$\$\$\$\$\$\$\$\$. The instantaneous oscillating electric field of this em wave is along +y axis. Then the direction of oscillating magnetic field of the em wave will be along
 - (1) + z direction
 - (2) y direction
 - (3) z direction
 - (4) -x direction
- 17. The refractive index of the material of a prism is $\sqrt{2}$ and the angle of the prism is 30° . One of the two refracting surfaces of the prism is made a mirror inwards, by silver coating. A beam of monochromatic light entering the prism from the other face will retrace its path (after reflection from the silvered surface) if its angle of incidence on the prism is
 - (1) 45°
 - (2) **30**°
 - (3) 60°
 - (4) zero
- 18. An object is placed at a distance of 40 cm from a concave mirror of focal length 15 cm. If the object is displaced through a distance of 20 cm towards the mirror, the displacement of the image will be
 - (1) 36 cm away from the mirror
 - (2) 30 cm towards the mirror
 - (3) 30 cm away from the mirror
 - (4) 36 cm towards the mirror
- 19. The magnetic potential energy stored in a certain inductor is 25 mJ, when the current in the inductor is 60 mA. This inductor is of inductance
 - (1) 138·88 H
 - $(2) \quad 1{\cdot}389 \ H$
 - $(3) \quad 0.138 \text{ H}$
 - (4) 13·89 H

An electron of mass m with an initial velocity $\overrightarrow{V} = V_0 \hat{i} \ (V_0 > 0)$ enters an electric field $\overrightarrow{E} = -E_0 \hat{i} \ (E_0 = \text{constant} > 0)$ at t = 0. If λ_0 is its de-Broglie wavelength initially, then its de-Broglie wavelength at time t is

- $\begin{array}{ll} (1) & \lambda_0 \left(1 + \frac{eE_0}{mV_0}t\right) \\ (2) & \lambda_0 t \\ (3) & \displaystyle \frac{\lambda_0}{\left(1 + \frac{eE_0}{mV_0}t\right)} \\ (4) & \lambda_0 \end{array}$
- **21.** For a radioactive material, half-life is 10 minutes. If initially there are 600 number of nuclei, the time taken (in minutes) for the disintegration of 450 nuclei is
 - (1) 10
 - (2) 30
 - (3) 20
 - (4) 15
- **22.** The ratio of kinetic energy to the total energy of an electron in a Bohr orbit of the hydrogen atom, is
 - (1) 1:-1
 - (2) 2:-1
 - (3) 1:1
 - (4) 1:-2
- **23.** When the light of frequency $2v_0$ (where v_0 is threshold frequency), is incident on a metal plate, the maximum velocity of electrons emitted is v_1 . When the frequency of the incident radiation is increased to $5v_0$, the maximum velocity of electrons emitted from the same plate is v_2 . The ratio of v_1 to v_2 is
 - (1) 1:4
 - (2) 4:1
 - (3) 1:2

2:1


(4)

- 24. Unpolarised light is incident from air on a plane surface of a material of refractive index 'μ'. At a particular angle of incidence 'i', it is found that the reflected and refracted rays are perpendicular to each other. Which of the following options is correct for this situation ?
 - (1) Reflected light is polarised with its electric vector perpendicular to the plane of incidence


(2)
$$i = \sin^{-1}\left(\frac{1}{\mu}\right)$$

- (3) Reflected light is polarised with its electric vector parallel to the plane of incidence
- $(4) \quad i = \tan^{-1}\left(\frac{1}{\mu}\right)$
- **25.** An astronomical refracting telescope will have large angular magnification and high angular resolution, when it has an objective lens of
 - (1) large focal length and small diameter
 - (2) large focal length and large diameter
 - (3) small focal length and large diameter
 - (4) small focal length and small diameter
- 26. In Young's double slit experiment the separation d between the slits is 2 mm, the wavelength λ of the light used is 5896 Å and distance D between the screen and slits is 100 cm. It is found that the angular width of the fringes is 0.20°. To increase the fringe angular width to 0.21° (with same λ and D) the separation between the slits needs to be changed to
 - (1) 1·9 mm
 - (2) $2 \cdot 1 \text{ mm}$
 - (3) 1·8 mm
 - $(4) \quad 1{\cdot}7\ mm$

In the circuit shown in the figure, the input voltage V_i is 20 V, $V_{BE} = 0$ and $V_{CE} = 0$. The values of I_B , I_C and β are given by



- (1) $I_B = 25 \ \mu A, \ I_C = 5 \ mA, \ \beta = 200$
- (2) $I_B = 20 \ \mu A, \ I_C = 5 \ mA, \ \beta = 250$
- (3) $I_B = 40 \ \mu A, \ I_C = 10 \ mA, \ \beta = 250$
- (4) $I_B = 40 \ \mu A, \ I_C = 5 \ mA, \ \beta = 125$
- **28.** In a p-n junction diode, change in temperature due to heating
 - (1) affects only forward resistance
 - (2) does not affect resistance of p-n junction
 - (3) affects only reverse resistance
- **29.** In the combination of the following gates the output Y can be written in terms of inputs A and B as



Downloadestreerer.com

- 30. A tuning fork is used to produce resonance in a glass tube. The length of the air column in this tube can be adjusted by a variable piston. At room temperature of 27°C two successive resonances are produced at 20 cm and 73 cm of column length. If the frequency of the tuning fork is 320 Hz, the velocity of sound in air at 27°C is
 - $(1) \quad 339 \text{ m/s}$
 - $(2) \quad 350 \text{ m/s}$
 - $(3) \quad \ \ 330 \ m/s$
 - $(4) \quad \ \ 300 \ m/s$
- **31.** The electrostatic force between the metal plates of an isolated parallel plate capacitor C having a charge Q and area A, is
 - (1) linearly proportional to the distance between the plates.
 - (2) proportional to the square root of the distance between the plates.
 - (3) independent of the distance between the plates.
 - (4) inversely proportional to the distance between the plates.
- **32.** An electron falls from rest through a vertical distance h in a uniform and vertically upward directed electric field E. The direction of electric field is now reversed, keeping its magnitude the same. A proton is allowed to fall from rest in it through the same vertical distance h. The time of fall of the electron, in comparison to the time of fall of the proton is
 - (1) 5 times greater
 - (2) 10 times greater
 - (3) smaller
 - (4) equal
- **33.** A pendulum is hung from the roof of a sufficiently high building and is moving freely to and fro like a simple harmonic oscillator. The acceleration of the bob of the pendulum is 20 m/s^2 at a distance of 5 m from the mean position. The time period of oscillation is
 - (1) πs
 - (2) 2 s
 - $(3) \quad 2\pi \; s$
 - (4) 1 s

- A metallic rod of mass per unit length 0.5 kg m^{-1} is lying horizontally on a smooth inclined plane which makes an angle of 30° with the horizontal. The rod is not allowed to slide down by flowing a current through it when a magnetic field of induction 0.25 T is acting on it in the vertical direction. The current flowing in the rod to keep it stationary is
- (1) 5.98 A
- (2) 14·76 A
- (3) 7·14 A
- (4) 11·32 A
- **35.** Current sensitivity of a moving coil galvanometer is 5 div/mA and its voltage sensitivity (angular deflection per unit voltage applied) is 20 div/V. The resistance of the galvanometer is
 - (1) 25 Ω
 - (2) 250 Ω
 - (3) 40 Ω
 - $(4) \quad 500 \ \Omega$
- **36.** A thin diamagnetic rod is placed vertically between the poles of an electromagnet. When the current in the electromagnet is switched on, then the diamagnetic rod is pushed up, out of the horizontal magnetic field. Hence the rod gains gravitational potential energy. The work required to do this comes from
 - (1) the magnetic field
 - (2) the lattice structure of the material of the rod
 - (3) the current source
 - (4) the induced electric field due to the changing magnetic field

37. An inductor 20 mH, a capacitor 100 μ F and a resistor 50 Ω are connected in series across a source of emf, V = 10 sin 314 t. The power loss in the circuit is

- $(1) \quad 0.43 \text{ W}$
- (2) 2.74 W
- $(3) \quad 0.79 \text{ W}$
- (4) 1.13 W


- The power radiated by a black body is P and it 42. 38. radiates maximum energy at wavelength, λ_0 . If the temperature of the black body is now changed so that it radiates maximum energy at wavelength $\frac{3}{4}\lambda_0$, the power radiated by it becomes nP. The value of n is
 - 4 (1) $\overline{3}$
 - 256(2)81
 - (3)81

(4)
$$\frac{61}{256}$$


- 39. Two wires are made of the same material and have the same volume. The first wire has cross-sectional area A and the second wire has cross-sectional area 3A. If the length of the first wire is increased by Δl on applying a force F. how much force is needed to stretch the second wire by the same amount?
 - (1)6 F
 - 4 F
 - (3)
 - (4)
- 40. A small sphere of radius 'r' falls from rest in a viscous liquid. As a result, heat is produced due **44**. to viscous force. The rate of production of heat when the sphere attains its terminal velocity, is proportional to
 - r^2 (1)
 - r^5 (2)
 - r^3 (3)
 - r^4 (4)
- A sample of 0.1 g of water at 100°C and normal $|_{45}$. 41. pressure $(1{\cdot}013~{\times}~10^5~Nm^{-2})$ requires 54 cal of heat energy to convert to steam at 100°C. If the volume of the steam produced is 167.1 cc, the change in internal energy of the sample, is
 - (1)208·7 J
 - (2) $42 \cdot 2 J$
 - (3)104·3 J 84·5 J (4)

- (2)
- 9 F
- F

- (1)D $\frac{7}{5}$ D (2)
 - Three objects, A : (a solid sphere), B : (a thin 43. circular disk) and C : (a circular ring), each have the same mass M and radius R. They all spin with the same angular speed ω about their own symmetry axes. The amounts of work (W) required to bring them to rest, would satisfy the relation
 - (1) $W_A > W_B > W_C$
 - $(2) \quad W_{\rm B} > W_{\rm A} > W_{\rm C}$
 - $(3) \quad W_{\rm C} > W_{\rm B} > W_{\rm A}$
 - $(4) \quad W_{A} > W_{C} > W_{B}$
 - A moving block having mass m, collides with another stationary block having mass 4m. The lighter block comes to rest after collision. When the initial velocity of the lighter block is v, then the value of coefficient of restitution (e) will be
 - 0.25(1)
 - (2)0.8
 - (3)0.5
 - (4)0.4
 - Which one of the following statements is incorrect?
 - (1)Limiting value of static friction is directly proportional to normal reaction.
 - (2)Frictional force opposes the relative motion.
 - (3)Rolling friction is smaller than sliding friction.
 - Coefficient (4)of sliding friction has dimensions of length.



46.	46. Match the metal ions given in Column I with the spin magnetic moments of the ions given in Column II and assign the <i>correct</i> code :						acid	A mixture of 2.3 g formic acid and 4.5 g oxalic acid is treated with conc. H_2SO_4 . The evolved gaseous mixture is passed through KOH pellets.		
		Colur	nn I		Column II		-	ght (in g) of the remaining product at STP		
	a.	C0 ³⁺		i.	$\sqrt{8}$ B.M.		will			
	b.	Cr^{3+}		ii.	$\sqrt{35}$ B.M.		(1) (2)	3·0 2·8		
	c.	Fe^{3+}		iii.	$\sqrt{3}$ B.M.		(2) (3)	1.4		
	d.	Ni ²⁺		iv.	$\sqrt{24}$ B.M.		(4)	4.4		
				v.	$\sqrt{15}$ B.M.	52.	The	difference between amylose and amylopectin		
		a	b	с	d		is			
	(1)	i	ii	iii	iv		(1)	Amylose have $1 \rightarrow 4$ α -linkage and $1 \rightarrow 6 \beta$ -linkage		
	(2) (3)	iv iv	i	ii ii	iii i		(2)	Amylopectin have $1 \rightarrow 4 \alpha$ -linkage and $1 \rightarrow 6 \beta$ -linkage		
	(4)	iii	v v	i	ü		(3)	Amylopectin have $1 \rightarrow 4$ α -linkage and $1 \rightarrow 6 \alpha$ -linkage		
47.	Iron	carbor	nyl, Fe($CO)_5$ is			(4)	Amylose is made up of glucose and		
	(1)		nuclear	r				galactose		
	(2)	trinu				53.	-	arding cross-linked or network polymers,		
	(3) (4)	dinuc	nuclear				whi (1)	ch of the following statements is <i>incorrect</i> ? They are formed from bi- and tri-functional		
10							(1)	monomers.		
48.		type Cl ₂ (en)		nerism s	shown by the complex		(2)	Examples are bakelite and melamine.		
	(1)		_	n isomeri	ism		(3)	They contain covalent bonds between various linear polymer chains.		
	(2)			omerism			(4)	They contain strong covalent bonds in their		
	(3)	Geom	netrical	isomeris	sm		(=)	polymer chains.		
	(4)	Linka	age ison	nerism		54.	Nitration of aniline in strong acidic medium			
49.					llowing ions exhibits agnetism as well?		-	s m-nitroaniline because		
			_	i parama	ignetism as well?		(1)	In electrophilic substitution reactions amino group is meta directive.		
	(1)	Cr_2O	•				(2)	In absence of substituents nitro group		
	(2)	MnO	$\frac{-}{4}$				(2)	always goes to m-position.		
	(3)	CrO_4^2	} ⊧				(3)	In spite of substituents nitro group always goes to only m-position.		
	(4)	MnO	$\frac{2}{4}$				(4)	In acidic (strong) medium aniline is present as anilinium ion.		
50.					netic behaviour of the	55.		ch of the following oxides is most acidic in		
			Ni(CO) ₄		1 1			ire ?		
	 (1) (2) 				y and diamagnetic		(1)	BeO BeO		
	(2) (3)	_	-	-	etry and paramagnetic etry and diamagnetic		(2) (3)	BaO MgO		
	(3) (4)	-	-	-	v and paramagnetic		(3) (4)	CaO		
	(-)			,,		I	(-)			



the electrophile involved is


- (1) formyl cation ($\overset{\smile}{CHO}$)
- (2) dichloromethyl anion $(CHCl_2)$
- (3) dichloromethyl cation (\breve{CHCl}_2)
- (4) dichlorocarbene ($:CCl_2$)
- **57.** Carboxylic acids have higher boiling points than aldehydes, ketones and even alcohols of comparable molecular mass. It is due to their
 - (1) formation of carboxylate ion
 - (2) more extensive association of carboxylic acid via van der Waals force of attraction
 - (3) formation of intramolecular H-bonding
 - (4) formation of intermolecular H-bonding
- 58. Compound A, $C_8H_{10}O$, is found to react with NaOI (produced by reacting Y with NaOH) and yields a yellow precipitate with characteristic smell.

A and Y are respectively

CH₃

(4)

- (2) (2) $CH CH_3 \text{ and } I_2$ I OH



 CH_3

 \rightarrow OH and I_2


- 59. The compound A on treatment with Na gives B, and with PCl₅ gives C. B and C react together to give diethyl ether. A, B and C are in the order
 - $(1) \quad C_2H_5OH, C_2H_5Cl, C_2H_5ONa$
 - (2) $C_2H_5Cl, C_2H_6, C_2H_5OH$
 - $(3) \quad \mathrm{C_2H_5OH,\,C_2H_6,\,C_2H_5Cl}$
 - (4) C_2H_5OH , C_2H_5ONa , C_2H_5Cl
- **60.** Which oxide of nitrogen is *not* a common pollutant introduced into the atmosphere both due to natural and human activity ?
 - (1) NO₂
 - (2) N₂O
 - (3) N₂O₅
 - (4) NO
- 61. Hydrocarbon (A) reacts with bromine by substitution to form an alkyl bromide which by Wurtz reaction is converted to gaseous hydrocarbon containing less than four carbon atoms. (A) is
 - (1) $CH_2 = CH_2$
 - $(2) \quad \mathrm{CH}_3 \mathrm{CH}_3$
 - $(3) \quad CH \equiv CH$
 - (4) CH₄
- **62.** The compound C_7H_8 undergoes the following reactions :


$$C_7H_8 \xrightarrow{3 \operatorname{Cl}_2/\Delta} A \xrightarrow{\operatorname{Br}_2/\operatorname{Fe}} B \xrightarrow{\operatorname{Zn}/\operatorname{HCl}} C$$

The product 'C' is

- (1) *o*-bromotoluene
- (2) 3-bromo-2,4,6-trichlorotoluene
- (3) *m*-bromotoluene
- (4) *p*-bromotoluene

63. Which of the following carbocations is expected to be most stable ?





- **64.** Which of the following is correct with respect to - I effect of the substituents ? (R = alkyl)
 - (1) $-NR_2 < -OR < -F$
 - (2) $-NH_2 > -OR > -F$
 - (3) $-NH_2 < -OR < -F$
 - $(4) \quad -\operatorname{NR}_2 > -\operatorname{OR} > -\operatorname{F}$
- **65.** Which of the following molecules represents the order of hybridisation sp², sp², sp, sp from left to right atoms ?
 - (1) $CH_2 = CH C \equiv CH$
 - (2) $CH_2 = CH CH = CH_2$
 - (3) $HC \equiv C C \equiv CH$
 - (4) $CH_3 CH = CH CH_3$

Identify the major products P, Q and R in the following sequence of reactions :

- **67.** Which of the following compounds can form a zwitterion ?
 - (1) Acetanilide
 - (2) Benzoic acid
 - (3) Aniline
 - (4) Glycine

68.	Following solutions were prepared by mixing different volumes of NaOH and HCl of different	72.	The bond dissociation energies of X_2 , Y_2 and XY are in the ratio of $1: 0.5: 1$. ΔH for the formation
	concentrations :		of XY is -200 kJ mol^{-1} . The bond dissociation
	a. 60 mL $\frac{M}{10}$ HCl + 40 mL $\frac{M}{10}$ NaOH		energy of X_2 will be
			(1) 100 kJ mol ⁻¹
	b. 55 mL $\frac{M}{10}$ HCl + 45 mL $\frac{M}{10}$ NaOH		(2) 800 kJ mol ⁻¹
	c. 75 mL $\frac{M}{5}$ HCl + 25 mL $\frac{M}{5}$ NaOH		(3) 200 kJ mol ^{-1}
	$\frac{1}{5}$		(4) 400 kJ mol ⁻¹
	d. 100 mL $\frac{M}{10}$ HCl + 100 mL $\frac{M}{10}$ NaOH pH of which one of them will be equal to 1 ?	73.	When initial concentration of the reactant is doubled, the half-life period of a zero order
	(1) a		reaction
	(2) d		(1) is doubled
	(3) b		(2) is tripled
	(4) c		(3) is halved
60	On which of the following properties does the		(4) remains unchanged
69.	On which of the following properties does the coagulating power of an ion depend ?	74.	For the redox reaction
	(1) Size of the ion alone		$\operatorname{MnO}_4^- + \operatorname{C}_2\operatorname{O}_4^{2-} + \operatorname{H}^+ \longrightarrow \operatorname{Mn}^{2+} + \operatorname{CO}_2 + \operatorname{H}_2\operatorname{O}$
	(2) Both magnitude and sign of the charge on the ion		the correct coefficients of the reactants for the balanced equation are
	(3) The magnitude of the charge on the ion alone		$MnO_4^ C_2O_4^{2-}$ H ⁺
	(4) The sign of charge on the ion alone		(1) 2 5 16
70.	Given van der Waals constant for NH_3 , H_2 , O_2		(2) 2 16 5
	and CO_2 are respectively 4.17, 0.244, 1.36 and		(3) 16 5 2
	3.59, which one of the following gases is most		(4) 5 16 2
	easily liquefied ?	75.	Which one of the following conditions will favour
	(1) H ₂		maximum formation of the product in the reaction,
	(2) O ₂		$A_2(g) + B_2(g) \rightleftharpoons X_2(g) \Delta_r H = -X kJ?$
	(3) NH ₃		 (1) Low temperature and low pressure
	(4) CO ₂		(1) How temperature and how pressure(2) High temperature and high pressure
71.	The solubility of $BaSO_4$ in water is		(3) Low temperature and high pressure
11.	2.42×10^{-3} gL ⁻¹ at 298 K. The value of its		(4) High temperature and low pressure
	solubility product (K_{sp}) will be	76.	The connection factor 's' to the ideal and equation
	(Given molar mass of $BaSO_4 = 233 \text{ g mol}^{-1}$)	70.	The correction factor 'a' to the ideal gas equation corresponds to
			(1) volume of the gas molecules
	(1) $1.08 \times 10^{-12} \text{ mol}^2 \text{ L}^{-2}$		(2) electric field present between the gas
	(2) $1.08 \times 10^{-14} \text{ mol}^2 \text{ L}^{-2}$		molecules
	(3) $1.08 \times 10^{-10} \text{ mol}^2 \text{ L}^{-2}$		(3) density of the gas molecules
	(4) $1.08 \times 10^{-8} \text{ mol}^2 \text{ L}^{-2}$		(4) forces of attraction between the gas molecules
	AC/KK/Page 11 Downloadest ACE FORM		

77.	The correct order of N-compounds in its decreasing order of oxidation states is	83.	The correct difference between first- and second-order reactions is that
	(1) HNO ₃ , NO, NH ₄ Cl, N ₂		(1) the half-life of a first-order reaction does not
	(2) HNO_3 , NH_4Cl , NO , N_2		depend on $[A]_0$; the half-life of a
	(3) HNO_3 , NO, N ₂ , NH ₄ Cl		 second-order reaction does depend on [A]₀ (2) a first-order reaction can be catalyzed; a
	(4) $\text{NH}_4\text{Cl}, \text{N}_2, \text{NO}, \text{HNO}_3$		(2) a first-order reaction can be catalyzed; a second-order reaction cannot be catalyzed
	(4) $1014_401, 102, 100, 11003_3$		(3) the rate of a first-order reaction does not
78.	Which one of the following elements is unable to $3-$		depend on reactant concentrations; the rate of a second-order reaction does depend on
	form MF_6^{3-} ion ?		reactant concentrations
	(1) Al		(4) the rate of a first-order reaction does
	(2) B		depend on reactant concentrations; the rate of a second-order reaction does not depend
	(3) Ga		on reactant concentrations
	(4) In	84.	Among CaH ₂ , BeH ₂ , BaH ₂ , the order of ionic
79.	Considering Ellingham diagram, which of the		character is
	following metals can be used to reduce alumina?		(1) $CaH_2 < BeH_2 < BaH_2$
	(1) Zn		(2) $BeH_2 < BaH_2 < CaH_2$
	(2) Mg (2) Fa		(3) $BeH_2 < CaH_2 < BaH_2$
	 (3) Fe (4) Cu 		(4) $BaH_2 < BeH_2 < CaH_2$
		85.	In which case is the number of molecules of water
80.	The correct order of atomic radii in group 13 elements is		maximum ?
	(1) $B < AI < Ga < In < TI$		(1) 0.18 g of water
	(2) $B < Ga < Al < Tl < In$		(2) 0.00224 L of water vapours at 1 atm and 273 K
	(3) B < Al < In < Ga < Tl		(3) 18 mL of water
	$(4) \mathbf{B} < \mathbf{G}\mathbf{a} < \mathbf{A}\mathbf{l} < \mathbf{I}\mathbf{n} < \mathbf{T}\mathbf{l}$		(4) 10^{-3} mol of water
81.	Which of the following statements is <i>not</i> true for	96	Consider the change in oxidation state of
01.	halogens ?	00.	Bromine corresponding to different emf values as
	(1) All are oxidizing agents.		shown in the diagram below :
	(2) All but fluorine show positive oxidation states.		$BrO_4^- \xrightarrow{1.82 \text{ V}} BrO_3^- \xrightarrow{1.5 \text{ V}} HBrO$
	(3) All form monobasic oxyacids.		$Br^{-} \leftarrow I \cdot 0652 V Br_{2} \leftarrow I \cdot 595 V$
	(4) Chlorine has the highest electron-gain enthalpy.		Br $\leftarrow 1.0652 \text{ V}$ Br ₂ $\leftarrow 1.595 \text{ V}$ Then the species undergoing disproportionation
82.	In the structure of ClF_3 , the number of lone pairs		is
	of electrons on central atom 'Cl' is		(1) $\operatorname{Br}O_{4}^{-}$
	(1) two		(2) Br_2
	(2) four		2
	(3) one(4) three		$\begin{array}{ccc} (3) & \operatorname{BrO}_3^- \\ (4) & \operatorname{HD}_2^- \end{array}$
			(4) HBrO

87. Consider the following species :

CN⁺, CN⁻, NO and CN

Which one of these will have the highest bond order?

- (1) CN⁻
- (2) CN⁺
- (3) NO
- (4) CN

88. Which one is a *wrong* statement ?

- (1) An orbital is designated by three quantum numbers while an electron in an atom is designated by four quantum numbers.
- $(2) \quad \ \ {\rm The \ electronic \ configuration \ of \ N \ atom \ is}$

$1s^2$	$2s^2$	$2p_x^1$	$2p_y^1$	$2p_z^1$
↑↓	↑↓	1	1	↓

- (3) Total orbital angular momentum of electron in 's' orbital is equal to zero.
- (4) The value of m for d_{z^2} is zero.
- **89.** Iron exhibits bcc structure at room temperature. Above 900°C, it transforms to fcc structure. The ratio of density of iron at room temperature to that at 900°C (assuming molar mass and atomic radii of iron remains constant with temperature) is
 - $(1) \quad \frac{4\sqrt{3}}{3\sqrt{2}}$
 - $(2) \quad \frac{3\sqrt{3}}{4\sqrt{2}}$
 - $(3) \quad \frac{\sqrt{3}}{\sqrt{2}}$
 - $(4) \quad \frac{1}{2}$

90. Magnesium reacts with an element (X) to form an ionic compound. If the ground state electronic configuration of (X) is 1s² 2s² 2p³, the simplest formula for this compound is

- (1) MgX₂
- (2) Mg₂X
- $(3) \quad Mg_2X_3$
- (4) Mg_3X_2

- **91.** Pollen grains can be stored for several years in liquid nitrogen having a temperature of
 - (1) $-80^{\circ}C$
 - $(2) 196^{\circ}C$
 - (3) $-120^{\circ}C$
 - $(4) 160^{\circ}C$
- 92. Oxygen is *not* produced during photosynthesis by
 - (1) Nostoc
 - (2) Cycas
 - (3) Green sulphur bacteria
 - (4) Chara
- **93.** What is the role of NAD⁺ in cellular respiration ?
 - (1) It functions as an electron carrier.
 - (2) It is a nucleotide source for ATP synthesis.
 - (3) It functions as an enzyme.
 - (4) It is the final electron acceptor for anaerobic respiration.
- **94.** Which of the following elements is responsible for maintaining turgor in cells ?
 - (1) Sodium
 - (2) Potassium
 - (3) Magnesium
 - (4) Calcium
- **95.** Which one of the following plants shows a very close relationship with a species of moth, where none of the two can complete its life cycle without the other ?
 - (1) Yucca
 - (2) Banana
 - (3) Hydrilla
 - (4) Viola
- **96.** In which of the following forms is iron absorbed by plants ?
 - (1) Ferrous
 - (2) Free element
 - (3) Ferric
 - (4) Both ferric and ferrous
- 97. Double fertilization is
 - (1) Fusion of one male gamete with two polar nuclei
 - $(2) \quad Fusion \ of \ two \ male \ gametes \ with \ one \ egg$
 - (3) Fusion of two male gametes of a pollen tube with two different eggs
 - (4) Syngamy and triple fusion

Downloadest4F6FARWaywww.workhapter.com

98.		ew' variety of rice was patented by a foreign	104.	Nicł	ne is
		pany, though such varieties have been		(1)	the physical space where an organism lives
	(1)	sent in India for a long time. This is related to Sharbati Sonora		(2)	the range of temperature that the organism
	(1) (2)	Lerma Rojo		$\langle \mathbf{O} \rangle$	needs to live
	(2)	Co-667		(3)	all the biological factors in the organism's environment
	(4)	Basmati		(4)	the functional role played by the organism
00				(=)	where it lives
99.		India, the organisation responsible for essing the safety of introducing genetically	105.	Whi	ch of the following is a secondary pollutant ?
		lified organisms for public use is		(1)	CO ₂
	(1)	Council for Scientific and Industrial		(2)	SO_2
		Research (CSIR)			-
	(2)	Research Committee on Genetic		(3)	СО
	(0)	Manipulation (RCGM)		(4)	O_3
	(3)	Indian Council of Medical Research (ICMR)	106.	Nata	ality refers to
	(4)	Genetic Engineering Appraisal Committee (GEAC)		(1)	Birth rate
				(2)	Number of individuals leaving the habitat
100.		ch of the following is commonly used as a or for introducing a DNA fragment in human		(3)	Death rate
		phocytes ?		(4)	Number of individuals entering a habitat
	(1)	Ti plasmid	107.	Wor	ld Ozone Day is celebrated on
	(2)	λ phage		(1)	21 st April
	(3)	Retrovirus		(2)	16 th September
	(4)	pBR 322			
101.		of bioresources by multinational companies		(3)	5 th June
		organisations without authorisation from the cerned country and its people is called		(4)	22 nd April
	(1)	Biopiracy	108.	Wha	at type of ecological pyramid would be
	(2)	Biodegradation		obta	ained with the following data ?
	(3)	Bio-infringement			Secondary consumer : 120 g
	(4)	Bioexploitation			Primary consumer : 60 g
109	The	correct order of stong in Polymorogy Chain			Primary producer : 10 g
102.		correct order of steps in Polymerase Chain ction (PCR) is		(1)	Pyramid of energy
	(1)	Annealing, Extension, Denaturation		(2)	Upright pyramid of numbers
	(2)	Denaturation, Extension, Annealing		(3)	Inverted pyramid of biomass
	(3)	Extension, Denaturation, Annealing		(4)	Upright pyramid of biomass
	(4)	Denaturation, Annealing, Extension	109.		tratosphere, which of the following elements
103.	Sele	ct the <i>correct</i> match :			as a catalyst in degradation of ozone and ase of molecular oxygen ?
	(1)	$F_2 \times \text{Recessive parent}$ – Dihybrid cross		(1)	Cl
	(2)	T.H. Morgan – Transduction		(1) (2)	Fe
	(3)	Ribozyme – Nucleic acid		(2)	Carbon
	(4)	G. Mendel – Transformation		(4)	Oxygen
	(-)		I	(1)	~~, <u>5</u> ~

HLAAC/KK/Page 14 Downloade@466FARvgvgH.@etchapter.com

110.		ch of the following pairs is <i>wrongly</i> ched?	117.		ondary xylem and phloem in dicot stem are luced by
	(1)	ABO blood grouping : Co-dominance		(1)	Vascular cambium
	(2)	XO type sex : Grasshopper		(2)	Phellogen
		determination		(3)	Apical meristems
	(3)	Starch synthesis in pea : Multiple alleles		(4)	Axillary meristems
	(4)	T.H. Morgan : Linkage	118		nts having little or no secondary growth are
111.	Sele	ct the <i>correct</i> statement :	110.	(1)	Deciduous angiosperms
	(1)	Punnett square was developed by a British		(1) (2)	Conifers
		scientist.		(2)	Grasses
	(2)	Spliceosomes take part in translation.		(4)	Cycads
	(3)	Franklin Stahl coined the term "linkage".			•
	(4)	Transduction was discovered by S. Altman.	119.		et potato is a modified
112.	The	experimental proof for semiconservative		(1)	Adventitious root
		ication of DNA was first shown in a		(2)	Tap root
	(1)	Bacterium		(3)	Stem
	(2)	Plant		(4)	Rhizome
	(3)	Fungus	120.	Pne	umatophores occur in
	(4)	Virus		(1)	Free-floating hydrophytes
113.	Sele	ct the <i>correct</i> match :		(2)	Carnivorous plants
	(1)	Alfred Hershey and – TMV		(3)	Halophytes
		Martha Chase		(4)	Submerged hydrophytes
	(2)	Matthew Meselson – Pisum sativum	121.	Casi	parian strips occur in
	(0)	and F. Stahl		(1)	Pericycle
	(3)	Alec Jeffreys – Streptococcus		(2)	Cortex
	(4)	<i>pneumoniae</i> Francois Jacob and – <i>Lac</i> operon		(3)	Epidermis
	(4)	Jacques Monod		(4)	Endodermis
114	Offa	-			
114.	(1)	ets are produced by Mitotic divisions	122.		ch of the following statements is <i>correct</i> ?
	(1) (2)	Parthenocarpy		(1)	Selaginella is heterosporous, while Salvinia
	(3)	Meiotic divisions		(\mathbf{O})	is homosporous.
	(4)	Parthenogenesis		(2)	Horsetails are gymnosperms.
115.	Whie	ch of the following flowers only once in its		(3)	Ovules are not enclosed by ovary wall in gymnosperms.
		time ?		(4)	Stems are usually unbranched in both
	(1)	Jackfruit		(4)	Cycas and Cedrus.
	(2)	Mango			-
	(3)	Bamboo species	123.		ct the <i>wrong</i> statement :
	(4)	Papaya		(1)	Mushrooms belong to Basidiomycetes.
116.		ch of the following has proved helpful in		(2)	Pseudopodia are locomotory and feeding
	-	erving pollen as fossils ? Cellulosic intine			structures in Sporozoans.
	(1) (2)	Oil content		(3)	Cell wall is present in members of Fungi and Plantae.
	(2) (3)	Pollenkitt		(A)	
	(3) (4)	Sporopollenin		(4)	Mitochondria are the powerhouse of the cell in all kingdoms except Monera.
<u> </u>		Sporoponenin	1		

124.	24. Match the items given in Column I with those in Column II and select the <i>correct</i> option given below :						suga	sugars are			
	0010	••• . Column	T		Column II			carbonyl and phosphate			
	0	Herbai		;	It is a place having a			hydroxyl and methyl			
	a.	Herbar	Tum	1.	collection of preserved		(4)	carbonyl and hydroxyl			
					plants and animals.	129.		ch of the following is not a product of light			
	b.	Key		ii.	A list that enumerates		react	tion of photosynthesis ?			
					methodically all the			NADH			
					species found in an area			NADPH			
					with brief description			ATP			
					aiding identification.	190		Oxygen			
	c.	Museu	m	iii.	Is a place where dried and	130.		ch among the following is <i>not</i> a prokaryote ? <i>Mycobacterium</i>			
					pressed plant specimens			Nostoc			
					mounted on sheets are			Saccharomyces			
	1				kept.			Oscillatoria			
	d.	Catalo	gue	1V.	A booklet containing a list of characters and their	131.		natal movement is <i>not</i> affected by			
					alternates which are			Light			
					helpful in identification o	f		O_2 concentration			
					various taxa.	-	(3)	Temperature			
		a	b	с	d			CO_2 concentration			
	(1)	iii	ii	i	iv	132.	The (Golgi complex participates in			
	(2)	ii	iv	iii	i		(1)	Formation of secretory vesicles			
	(3)	i	iv	iii	ii		(2)	Respiration in bacteria			
	(4)	iii	iv	i	ii		(3)	Fatty acid breakdown			
195	۸.ft-a			. falla	wed by maioria anaroa an		(4)	Activation of amino acid			
120.		г кагуо _§ luced ex			wed by meiosis, spores are	133.	Which of the following is true for nucleolus ?				
	(1)	Alterno	U	ousiy			(1)	It is a membrane-bound structure.			
		Agaric					(2)	It takes part in spindle formation.			
	(3)	Neuros		ţ			(3)	Larger nucleoli are present in dividing cells.			
	(4)	Saccha	-				(4)	It is a site for active ribosomal RNA			
100								synthesis.			
126.			en gr	ains a	re present in	134.		stage during which separation of the paired			
	(1)	Cycas Mongo						ologous chromosomes begins is			
	(2) (3)	Mango Mustar					 (1) Diplotene (2) Diakinesis 				
	(3) (4)	Pinus	u				(2) Diakinesis(3) Pachytene				
								Zygotene			
127.					matched ?	135.		nata in grass leaf are			
	(1)	-		-	oores – Brown algae			Kidney shaped			
	(2)	Gemm	-		– Marchantia			Rectangular			
	(3)				netes – Polysiphonia			Dumb-bell shaped			
_	(4)	Unicel	lular	organ	nism – <i>Chlorella</i>		(4)	Barrel shaped			

			1
136.		al bodies are mainly composed of	142
	(1)	DNA and RNA	
	(2)	Nucleic acids and SER	
	(3)	1	
	(4)	Free ribosomes and RER	
137.	Whi	ch of these statements is <i>incorrect</i> ?	143
	(1)	Glycolysis occurs in cytosol.	
	(2)	Glycolysis operates as long as it is supplied with NAD that can pick up hydrogen atoms.	
	(3)	Enzymes of TCA cycle are present in mitochondrial matrix.	
	(4)	Oxidative phosphorylation takes place in outer mitochondrial membrane.	
138.		ch of the following terms describe human tition?	144
	(1)	Thecodont, Diphyodont, Heterodont	
	(2)	Pleurodont, Monophyodont, Homodont	
	(3)	Thecodont, Diphyodont, Homodont	
	(4)	Pleurodont, Diphyodont, Heterodont	
139.	Sele	ct the <i>incorrect</i> match :	145
	(1)	Allosomes – Sex chromosomes	
	(2)	Submetacentric – L-shaped chromososmes chromosomes	
	(3)	Lampbrush – Diplotene bivalents chromosomes	
	(4)	Polytene – Oocytes of amphibians chromosomes	146
140.		ch of the following events does <i>not</i> occur in gh endoplasmic reticulum ?	
	(1)	Protein glycosylation	
	(2)	Cleavage of signal peptide	
	(3)	Protein folding	
	(4)	Phospholipid synthesis	
141.	mRI sim	ny ribosomes may associate with a single NA to form multiple copies of a polypeptide ultaneously. Such strings of ribosomes are ned as	
	(1)	Polyhedral bodies	
	(2)	Plastidome	
			1

- **142.** All of the following are part of an operon *except*
 - (1) structural genes
 - (2) an enhancer
 - (3) an operator
 - (4) a promoter
- 143. A woman has an X-linked condition on one of her X chromosomes. This chromosome can be inherited by
 - (1) Only sons
 - (2) Only grandchildren
 - (3) Only daughters
 - (4) Both sons and daughters
- **144.** According to Hugo de Vries, the mechanism of evolution is
 - (1) Saltation
 - (2) Phenotypic variations
 - (3) Multiple step mutations
 - (4) Minor mutations
- **145.** AGGTATCGCAT is a sequence from the coding strand of a gene. What will be the corresponding sequence of the transcribed mRNA ?
 - (1) UGGTUTCGCAT
 - (2) ACCUAUGCGAU
 - (3) AGGUAUCGCAU
 - (4) UCCAUAGCGUA
- 146. Match the items given in Column I with those in Column II and select the *correct* option given below :

	Colum	n I		Column II				
a.	Prolife	erative F	Phase	i.	Breakdown of endometrial lining			
b.	Secretory Phase				Follicular Phase			
c.	Menst	ruation		iii.	Luteal Phase			
	a	b	с					
(1)	i	iii	ii					
(2)	ii	iii	i					
(3)	iii	ii	i					
(4)	iii	i	ii					

Polysome

Nucleosome

(3)

(4)

147.		ch one of the following population ractions is widely used in medical science for	152.	Among the following sets of examples for divergent evolution, select the <i>incorrect</i> option :
		production of antibiotics ?		(1) Heart of bat, man and cheetah
	(1)	Mutualism		(2) Brain of bat, man and cheetah
	(2)	Parasitism		(3) Forelimbs of man, bat and cheetah
	(3)	Commensalism		(4) Eye of octopus, bat and man
	(4)	Amensalism	153.	
148	A11	of the following are included in 'Ex-situ	100.	disease ?
140.		servation' <i>except</i>		(1) Rheumatoid arthritis
	(1)	Sacred groves		(2) Alzheimer's disease
	(2)	Botanical gardens		(3) Psoriasis
	(3)	Wildlife safari parks		(4) Vitiligo
	(4)	Seed banks	154.	In which disease does mosquito transmitted
149.		ch the items given in Column I with those in umn II and select the <i>correct</i> option given		pathogen cause chronic inflammation of lymphatic vessels ? (1) Ascariasis
	Dero	Column I Column II		(2) Ringworm disease
	a.	Eutrophication i. UV-B radiation		(3) Elephantiasis
	a. b.	Sanitary landfill ii. Deforestation		(4) Amoebiasis
	с.	Snow blindness iii. Nutrient	155	Conversion of milk to curd improves its
		enrichment	100.	nutritional value by increasing the amount of
	d.	Jhum cultivation iv. Waste disposal		(1) Vitamin A
		a b c d		(2) Vitamin B ₁₂
	(1)	i iii iv ii		(3) Vitamin D
	(2)	iii iv i ii		(4) Vitamin E
	(3)	ii i iii iv	156	The similarity of bone structure in the forelimbs
	(4)	i ii iv iii	100.	of many vertebrates is an example of
150	Ing	growing population of a country,		(1) Analogy
100.		reproductive individuals are less than the		(2) Convergent evolution
	(1)	post-reproductive individuals are less than the		(3) Homology
	(2)	reproductive and pre-reproductive		(4) Adaptive radiation
		individuals are equal in number.	157.	8 1
	(3)	pre-reproductive individuals are more than		'Inheritance of blood groups' in humans ? a. Dominance
	(\mathbf{A})	the reproductive individuals.		b. Co-dominance
	(4)	pre-reproductive individuals are less than the reproductive individuals.		c. Multiple allele
		-		d. Incomplete dominance
151.		ch part of poppy plant is used to obtain the		e. Polygenic inheritance
		g "Smack" ?		(1) a, b and c
	(1) (2)	Latex Roots		(2) b, d and e
	(2) (3)	Flowers		(3) b, c and e
	(3) (4)	Leaves		(4) a, c and e
	(4)	LEAVES		

158.	Horr	nones secreted by the placenta to maintain	162.	Whi	ch of the following is an	amino acid derived	
	preg	mancy are		hormone ?			
	(1)	hCG, hPL, estrogens, relaxin, oxytocin		(1)	Ecdysone		
	(2)	hCG, hPL, progestogens, estrogens		(2)	Estradiol		
	(3)	hCG, hPL, progestogens, prolactin		(3)	Epinephrine		
	(4)	hCG, progestogens, estrogens, glucocorticoids		(4)	Estriol		
159.	The	contraceptive 'SAHELI'	163.		ch of the following stru prrectly paired with its f	-	
	(1)	increases the concentration of estrogen and prevents ovulation in females.		(1)	Limbic system :	consists of fibre tracts that	
	(2)	is an IUD.				interconnect	
	(3)	blocks estrogen receptors in the uterus, preventing eggs from getting implanted.				different regions of brain; controls movement.	
	(4)	is a post-coital contraceptive.		(2)	Hypothalamus :	production of	
160.	from					releasing hormones and regulation of temperature, hunger and thirst.	
	(1)	endoderm and mesoderm		(3)		controls respiration	
	(2)	mesoderm and trophoblast		(0)	-	and cardiovascular	
	(3) (4)	ectoderm and mesoderm ectoderm and endoderm				reflexes.	
161.		difference between spermiogenesis and		(4)	-	band of fibers connecting left and right cerebral	
	sper	miation is				hemispheres.	
	(1)	In spermiogenesis spermatozoa are formed, while in spermiation spermatids are formed.			transparent lens in the l lace by	human eye is held in	
	(2)	In spermiogenesis spermatozoa from sertoli		(1)	ligaments attached to t		
		cells are released into the cavity of		(2)	smooth muscles attache		
		seminiferous tubules, while in spermiation		(3)	с , , , , , , , , , , , , , , , , , , ,		
		spermatozoa are formed.		(4)	smooth muscles attache	ed to the ciliary body	
	(3)	In spermiogenesis spermatids are formed, while in spermiation spermatozoa are formed.	165.		ch of the following ho ificant role in osteoporos		
	(4)	In spermiogenesis spermatozoa are formed,		(1)	Progesterone and Aldos		
		while in spermiation spermatozoa are		(2)	Estrogen and Parathyr		
		released from sertoli cells into the cavity of		(3)	Aldosterone and Prolac		
		seminiferous tubules.		(4)	Parathyroid hormone a	na Prolactin	

166.	Which of the following options correctly represents the lung conditions in asthma and emphysema, respectively ?						169.	59. Which of the following gastric cells indirectly help in erythropoiesis ?					ectly
	(1)	(1) Increased number of bronchioles; Increased respiratory surface						(1)					
	(2)	Increased respiratory surface; Inflammation of bronchioles						(2) Goblet cells(3) Chief cells					
	(3)	Inflammation of bronchioles; Decreased respiratory surface						(4)					
	(4)	Decreased respiratory surface; Inflammation of bronchioles					170.	0. Match the items given in Column I with those in Column II and select the <i>correct</i> option given					
167.	Match the items given in Column I with those in Column II and select the <i>correct</i> option given							below :					5- 1 0
		below :						Column I				Column II	
		Column I Column II				Column II		a.	Fibriı	nogen	i.	Osmotic balance	
	a.	Tricuspid valve i.				Between left atrium		b.	Globu	ılin	ii.	Blood clotting	
	b.	Bicus	pid valv	re ii		and left ventricle Between right		c.	Albur	nin	iii.	Defence mechani	ism
		-	-			ntricle and			a	b	с		
		a			-	lmonary artery		(1)	i	ii	iii		
	c.					tween right rium and right		(2)	i	iii	ii		
						ventricle		(3)	iii	ii	i		
		a	b	С				(4)	ii	iii	i		
	(1)	i	iii	ii			171	Whi	eh of	the	following	is an occupat	ional
	(2)	i	ii	iii			171.		oiratory		-	is all occupat	101141
	(3)	iii 	i	ii 				-					
	(4)	ii	i	iii				(1) Silicosis					
168.		Match the items given in Column I with those in							(2) Botulism				
	Column II and select the <i>correct</i> option given below :							(3)	(3) Anthracis				
		Column I				Column II		(4) Emphysema					
	a.	Tidal volume			i.	i. 2500 – 3000 mL	172.	Calcium is important in skeletal muscle					
	b.	Inspiratory Reserve ii. 1100 – 1200 mL volume						cont	contraction because it				
								(1)				ng to	
	c.	Expiratory Reserve iii. 500 – 550 mL volume							it.				
	d.	Residual volume iv. 1000 – 1100 mL						(2) detaches the myosin head from the actin			actin		
		a b c		d	_			filam					
	(1)	iii	i	iv	ii			(3) binds to troponin to remove the masking of active sites on actin for myosin.				ng of	
	(2)	i	iv	ii	iii			(4)				ion of bonds bet	woon
	(3)	iii	ii	i	iv			(1)	-			ridges and the	
	(4)	iv	iii	ii	i				filam				
			00	Dei		odost/estop							nalich

173.	 Match the items given in Column I with the Column II and select the <i>correct</i> option below : 						Which of the following features is used to identify a male cockroach from a female cockroach ?(1) Presence of caudal styles			
		a. Glycosuria i. Acc		Column II		(2) (3)	Presence of a boat shaped sternum on the 9 th abdominal segment			
	a.			Accumulation of uric acid in joints		(4)				
	b.	Gout Renal calculi		ii.	Mass of crystallised salts within the kidne	y 176.	Identify the vertebrate group of animals characterized by crop and gizzard in its digestive system.(1) Reptilia			
	c.			iii.	Inflammation in glomeruli					
	d. Glomerular nephritis		iv.	Presence of glucose in urine		(2)	•			
		a	ı b c		d		(4)	Osteichthyes		
	(1) (2)	i ii	ii iii	iii i	iv iv	177.	Which one of these animals is not a homeotherm?			
	(3)	iii	ii	iv	i		(1)	Chelone		
	(4)	iv	i	ii	iii		(2)	Camelus		
174.	Match the items given in Column I with those in Column II and select the <i>correct</i> option given below :						(3) Macropus			
							(4) Psittacula			
		Column I (Function)			Column II	178.	Which of the following organisms are known as chief producers in the oceans ?			
					(Part of Excretory System)	,	(1) Diatoms			
	a.	. Ultrafiltration		n	i. Henle's loop		(2) Cyanobacteria			
		Concentration of urine			-		(3)	0		
	b.				ii. Ureter		(4)	Euglenoids		
	c.	Trans urine	port of		iii. Urinary bladder	179.	Which of the following animals does <i>not</i> undergo metamorphosis ?			
	d.	Storage of urine			iv. Malpighian corpuscle		(1) Tunicate			
							(2) (3)	Moth Earthworm		
					v. Proximal convoluted tubul		(4)	Starfish		
			h		d		Cilia	ates differ from all other protozoans in		
	(1)	a iv	b i	с ii	a iii		(1)	having a contractile vacuole for removing		
	(1)	v	iv	ii	ii			excess water		
	(2)	v iv	v	ii	iii		(2) (3)	using pseudopodia for capturing prey using flagella for locomotion		
			v iv	ii	iii		(3)	having two types of nuclei		
	(4)	v	10	1	111		(- /			

SPACE FOR ROUGH WORK

SPACE FOR ROUGH WORK

Read carefully the following instructions :

- 1. Each candidate must show on demand his/her Admit Card to the Invigilator.
- 2. No candidate, without special permission of the Superintendent or Invigilator, would leave his/her seat.
- 3. The candidates should not leave the Examination Hall without handing over their Answer Sheet to the Invigilator on duty and sign the Attendance Sheet twice. **Cases where a** candidate has not signed the Attendance Sheet second time will be deemed not to have handed over the Answer Sheet and dealt with as an unfair means case.
- 4. Use of Electronic/Manual Calculator is prohibited.
- 5. The candidates are governed by all Rules and Regulations of the examination with regard to their conduct in the Examination Hall. All cases of unfair means will be dealt with as per Rules and Regulations of this examination.
- 6. No part of the Test Booklet and Answer Sheet shall be detached under any circumstances.
- 7. The candidates will write the Correct Test Booklet Code as given in the Test Booklet/Answer Sheet in the Attendance Sheet.