Name of Student:	Roll no.
Pre-Board Assessmen	
	SS: 12
Time allowed: 3 hours	Chemistry Maximum Marks: 70
General Instructions	
 This question paper contains 35 quest In SECTION-A: Question no. 1 to 18 carrying 1 mark each. 	
➤ In SECTION-B: Question no. 19 to 25 (VSA) carrying 2 marks each.	are very short answer type questions
In SECTION-C: Question no. 26 to 30 carrying 3 marks each.	are short answer type questions (SA)
In SECTION-D: Question no. 31 and marks each.	32 are case based question carrying 4
In SECTION-E: Question no. 33 to 35 carrying 5 marks each.	are long answer type questions (LA)
SECTION-A (OBJECTIVE	TYPE QUESTIONS)
Each question c	
1) The values of van't hoff factor for KCI,	
a) 2,2,2	c) 1,1,1
b) 2,2,3	d) 1,1,2
2) Standard Electrode Potential for Standard	
a) - 0.5V b) +2.0V 3) The role of catalyst is to change	c) + 1.0V d) 0.0V
a) Enthalpy of reaction	c) Gibbs free energy of reaction
b) Activation energy of reaction4) The stability of ferric ion is due to	d) All the of above
a) Half-filled d- orbitals	c) Half-filled f- orbitals
b) completely filled f- orbitals	d) completely filled d- orbitals

. I with copper al	573K gives	
5) Dehydration of secondary alcohol with copper at	c) aldehyde d) ether	
a) alkenes b) ketone 6) Molecules whose mirror is non-superimposable or	ver them are known as chiral.	
6) Molecules whose mirror is non-superimposable of	Ver une	
Which of the following mirror images is chiral?	c) 1-bromobutane	
a) 2- bromobutane	c) 1-biomosatz	
b) 2-bromopropane	d) 2-bromopropan-2-ol	
7) Glycogen is an example of		
a) monosaccharide's	e) polysaccharide's	
h) Disambarida's	d) vitamins	
8) When attraction between A-B is less than that of	AA and BB, the solution will	
show which type of deviation from Raoult's law		
a) negative b) positive c) both	d) no deviation	
9) Aldehyde and ketone can be distinguished by		
√a) NaHSO₃	c) Silver nitrate	
b) Fehling solution	d) concentrated sulphuric acid	
10) The units of molar conductivity is:		
∠a) S m²mol⁻¹	c) S Cm ² mol ⁻¹	
b) S m ² mol ¹	d) S m ³ mol ⁻¹	
11) The IUPAC name of (CH ₃) ₂ -C=CH-COCH ₃		
a) 2-Methylpent-3-ene-4-one	c) 1,1-Dimethyl-2-ene-1-one	
b) 4-Methylpent-3-ene-2-one	d) 2,4-Dimethyl-2-ene-4-one	
12) The value of Henry's constant-		
a) increases with increase in temperature		
b) decreases with increase in temperature		
c) remains constant		
,d) first increases then decreases		
13) The atomic number of cerium is 58. The correct	ct electronic configuration of Ce3+	
is:		
a) [Xe]4f1 b) [Kr]4f1 c) [Xe]4f13	d) [Kr]4d ¹	
14) Identify the order of reaction when units of rate		
a) Second b) Zero c) Third	d) First	
(15-18) Assertion - Reason based questions:		
These question consists of two statements each printed as Assertion and Reason.		
While answering these questions you are required to choose any one of the		

Page 2 of 6

allowing responses xplanation of the 3 not correct es alse. D. If bo

alse. U.SER

o) ether

following responses. A. If both Assertion and Reason are true, Reason is correct explanation of the Assertion. B. If both Assertion and Reason are true but Reason is not correct explanation of the Assertion. C. If Assertion is true but Reason is false. D. If both Assertion and Reason are false

- 15) ASSERTION (A): Molality of the solution does not change with change in temp.

 REASON(R) : The molarity is expressed as number of moles per 1000g of the solvent.
- 16) ASSERTION (A): Oxidation state of Ni in [Ni(CN)4]2- is +2.
- REASON(R): The charge on cyanide ion is -1 and the total charge on the complex anion is -2.
- 17) ASSERTION (A): All aldehydes do not take part in aldol condensation.

 REASON(R): Carbanion is generated by the removal of alpha hydrogen by the base in the Aldol condensation.
- 18) ASSERTION (A): All amino acids exist as Zwitter ion.
 REASON (R): alpha- amino acids have both -NH₂ and -COOH group.

SECTION-B (VERY SHORT ANSWER TYPE QUESTIONS) Each question carries 2 marks.

- 19) Define Kohlrausch law. How does it help in calculation of molar conductivity at infinite dilution for a weak electrolyte CH₃COOH ?
- 20) Give two differences between DNA and RNA?
- 21) Time required to decompose thionyl chloride to half of its initial amount is 60 minutes. If the decomposition is a first order reaction, calculate the rate constant of the reaction.
- 22) Write equation for the preparation of 1- bromobutane from the following:
 - a) Butan-1-ol

- b) 1- chlorobutane
- 23) a) How will you prepare phenol from cumene?
 - b) Arrange the following in increasing order of acidic strength and Give reason.

 Phenol, p-nitrophenol, p-methyl phenol
- 24) Convert benzene diazonium chloride into:
 - a) Phénol

b) lodobenzene

OR

How p-aminoazobenzene can be obtained from aniline?

2, - Chi - Chi - Chi, - Chi;

entities?

b) [Co(NH₃)₃Cl₃]

a) [Cr(C2O4)3] -3

SECTION-C (SHORT ANSWER TYPE QUESTIONS)

Each question carries 3 marks.

- 26) Define isotonic solution. A three percent solution of sucrose C₁₂H₂₂O₁₁ is isotonic with 4% solution of an unknown organic compound. Calculate molecular mass of the unknown compound?
- 27) [Co (NH₃)₆]³⁺ is an inner orbital complex, whereas [Ni (NH₃)₆]²⁺ is an outer orbital complex?
- 28) Explain the following reactions:
 - a) Stephen Reduction
- b) Cannizzaro reduction
- c) Reimer Tiemann reaction

OR

Convert

- i) propanone to propene
- ii) ethanol to 3-hydroxybutanal
- iii) benzoic acid to benzaldehyde
- 29) Account for the following:
 - a) pKb of aniline is more than methylamine.
 - b) Ethylamine is soluble in water whereas aniline is not.
 - c) Gabriel Phthalimide Synthesis is preferred for synthesizing primary amines.
- 30) How are vitamins classified. Name the vitamin responsible for coagulation of blood?

SECTION D: CASE STUDY QUESTIONS.

Each question carries 4 marks.

Read the following passage carefully and answer the following questions:

31. Transition elements have partly filled d-orbitals in their normal oxidation states or in their common oxidation states. There are four series in transition elements and these are 3d- series (first transition series), 4d-series (second transition series), 5d-series (third transition series) and 6d-series (fourth transition series). Each series contains ten elements. Elements of group 12 are not considered as transition elements as these have fully-filled d-subshells. Transition elements show all the characteristics of metals. These elements show variable oxidation states, form

Page 4 of 6

coloured ions, form complexes from alloys and interstitial compounds. They show high enthalpies of atomisation, show catalytic properties and magnetic moment.

- (a) Why Zn, Cd and Hg are not considered as transition elements?
 - (b) Which element of d-block shows maximum oxidation state?
 - (c) Why transition elements show variable oxidation states?

Or

Why transition elements show higher enthalpies of atomization?

- (d) Calculate the magnetic moment of a divalent ion in its aqueous solution if its atomic number is 25.
- 32. In case of optically active alkyl halides, product formed as a result of $S_N 2$ mechanism has the inverted configuration as compared to the reactant? This is because the nucleophile attaches itself on the side opposite to the one where halogen atom is present thus, $S_N 2$ reactions of optically active halides are accompanied by inversion of configuration. In case of optically active alkyl halides $S_N 1$ reactions are accompanied by racemization. Actually the carbocation formed in the slow step being sp^2 hybridized is achiral. The attack of nucleophile is accomplished from either side resulting in mixture of products having same configuration.

Read the following passage carefully and answer the following questions:

- i) Define racemization. (1)
- ii) Write two differences between S_N1 and S_N2 reactions. (2)
- iii) Which alkyl halide from the following pair would react more rapidly by S_N2 mechanism, explain (1)

CH₃CH₂CH₂Br OR CH₃CH₂CH(Br)CH₃

OR

Which alkyl halide from the following pair would react more rapidly by S_N1 mechanism, explain

CH₃CH₂CH₂CH₂Br OR CH₃CH₂CH(Br)CH₃

SECTION E: LONG ANSWER TYPE QUESTIONS Each question carries 5 marks.

33) a) The rate constants of a reaction at 500 K and 700 K are 0.02 s⁻¹ and 0.07 s⁻¹ respectively.

Calculate the value of E_a (3)

	(2)
b) What are the factors affecting rate of reaction. OR	
	(2)
a) Define pseudo first order reaction with example? b) Define energy of activation. The rate of a particular reaction of a particular reaction.	doubles
b) Define energy of activation. The rate of a particular the when temperature changes from 27°C to 37°C. Calculate the	energy of
	(3)
activation?	(2)
34) a) Write equations of the following reactions:	
Sandmeyer reaction	
ii) Swart's reaction	(2)
b) Haloarenes are less reactive than haloalkanes why?	(2)
c) Write one use of DDT?	(1)
OR	
a) Primary alkyl halide C ₄ H ₉ Br (a) reacted with alcoholic KOH to give	compoun
(b). Compound (b) is reacted with HBr to give (c) which is an iso	omer of (a)
When (a) is reacted with sodium metal it gives compound (d), C ₈ H	18 which i
different from the compound formed when n-butyl bromide is reacted w	vith sodium
Give the structural formula of (a) and write the eq. for all the reactions.	(3)
b) How the following conversion can be carried out?	(2)
(i) Aniline into chlorobenzene	
(ii) Benzene into 4-Bromonitrobenzene	
35) a) Give the working of dry cell.	(2)
b) Represent the cell in which the following reaction takes place	(3)
$Mg (s) + 2Ag^{+} (0.0001M) \longrightarrow Mg^{2+} (0.130M) +$	2Ag (s)
Calculate its E _{cell} if E ^o _{cell} = 3.17 V.	
OR	
a) Give the working of fuel cell.	(2)
b) Calculate equilibrium constant for the following reaction	(3)
$Cu (s) + 2Ag^{+} (aq)$ \longrightarrow $2Ag (s) + Cu^{2+} (aq)$	
Given that $E^{o}_{Ag}^{+}/_{Ag} = 0.80V$ and $E^{o}_{cu}^{+2}/_{cu} = 0.34V$	
	Page 6 of 6